arXiv:math/0702022v1 [math.AP] 1 Feb 2007

SCATTERING POLES NEAR THE REAL AXIS FOR TWO STRICTLY CONVEX OBSTACLES.

ALEXEI IANTCHENKO

Abstract. To study the location of poles for the acoustic scattering matrix for two strictly convex obstacles with smooth boundaries, one uses an approximation of the quantized billiard operator M along the trapped ray between the two obstacles. Using this method Gérard (*cf.* [9]) obtained complete asymptotic expansions for the poles in a strip $\operatorname{Im} z \leq c$ as $\operatorname{Re} z$ tends to infinity. He established the existence of parallel rows of poles close to $\frac{\pi k}{d} + ij\delta$, $k \in \mathbb{Z}$, $j \in \mathbb{Z}_+$. Assuming that the boundaries are analytic and the eigenvalues of Poincaré map are non-resonant we use the Birkhoff normal form for M to improve his result and to get the complete asymptotic expansions for the poles in any logarithmic neighborhood of real axis.

1. Some known results on the localization of resonances.

1.1. Introduction.

Following [26] we denote by $-\Delta_{\mathbb{R}^{n+1}\setminus\mathcal{O}}$ the Dirichlet Laplacian on a connected exterior domain $\mathbb{R}^{n+1}\setminus\mathcal{O}$, where $\mathcal{O}$ is compact with a C^∞ boundary. Then the resolvent

$$R_{\mathcal{O}}(\lambda) \stackrel{\text{def}}{=} \left(-\Delta_{\mathbb{R}^{n+1}\setminus\mathcal{O}} - \lambda^2\right)^{-1} :\ L^2(\mathbb{R}^{n+1}\setminus\mathcal{O}) \mapsto H^2(\mathbb{R}^{n+1}\setminus\mathcal{O}) \cap H^1_0(\mathbb{R}^{n+1}\setminus\mathcal{O}),\ \operatorname{Im}\lambda < 0,$$

continues meromorphically across continuous spectrum $\operatorname{Im}\lambda = 0$, to an operator

$$R_{\mathcal{O}}(\lambda) :\ L^2_{\text{comp}}(\mathbb{R}^{n+1}\setminus\mathcal{O}) \mapsto H^2_{\text{loc}}(\mathbb{R}^{n+1}\setminus\mathcal{O}) \cap H^1_{0,\text{loc}}(\mathbb{R}^{n+1}\setminus\mathcal{O}).$$

Here $H^2(\mathbb{R}^{n+1}\setminus\mathcal{O})$ is the standard Sobolev space, $H^1_0(\mathbb{R}^{n+1}\setminus\mathcal{O})$ is the closure of $C^\infty_{\text{comp}}(\mathbb{R}^{n+1}\setminus\mathcal{O})$ in the H^1-norm and by L^2_{comp}, where by L^2_{comp} we mean the elements from L^2 that are zero outside some bounded set, and by H^2_{loc}, $H^1_{0,\text{loc}}$ functions that are locally in these spaces.

We recall that $R_{\mathcal{O}}(\lambda)$ is globally meromorphic in $\lambda \in \mathbb{C}$ when $n+1$ is odd and in $\lambda \in \Lambda$, the logarithmic covering of the complex plane, when $n+1$ is even.

Definition 1. *The poles of $R_{\mathcal{O}}$ are called resonances or scattering poles.*

Note that we adopted a convention that the resonances λ_j are in the upper half plane $\operatorname{Im}\lambda > 0$.

We cite some results following the review by Zworski [28] (note his convention that $\operatorname{Im}\lambda_j < 0$.)

An obstacle $\mathcal{O}$ contained in a ball $B(0,R)$ is called *non-trapping* if there exists $T > 0$ such that every broken characteristic ray (see e.g. [11], Section 24.2) starting in $(\mathbb{R}^{n+1}\setminus\mathcal{O}) \cap B(0,R)$ leaves that set within time T.

Propagation of singularities for boundary value problems, established by Melrose-Sjöstrand and Ivrii (see Chapter 24 of [11] and the references given), gives the basic results:

$$\begin{aligned} &\mathcal{O} \text{ is non-trapping, } \partial\mathcal{O} \text{ is } C^\infty \ \Rightarrow \\ (1)\qquad &\text{there exist } a, b > 0 \text{ such that a region } \{z;\ \operatorname{Im} z \leq a\log(1+|z|) + b\} \\ &\text{contains no poles;} \\ (2)\qquad &\forall N,\ \sharp\{\lambda_j :\ |\operatorname{Im}\lambda_j| \leq N\log(1+|\lambda_j|)\} < \infty. \end{aligned}$$

Date: February 2, 2008.

The situation is somewhat more satisfying in the analytic case where one can apply the propagation of Gevrey 3 singularities due to Lebeau [16]. In fact, Bardos-Lebeau-Rauch [2] (and also independently Popov [19]) used the work of Lebeau to show that

$$\begin{aligned} &\mathcal{O} \text{ is non-trapping, } \partial\mathcal{O} \text{ is real analytic } \Rightarrow \\ &\text{there exist } a, b > 0 \text{ such that a region } \{z;\ \operatorname{Im} z \leq a(1+|z|)^{1/3} + b\} \\ &\text{contains no poles.} \end{aligned} \tag{3}$$

Then clearly,

$$\exists B, C > 0:\ \sharp\{\lambda_j:\ |\operatorname{Im}\lambda_j| \leq B|\operatorname{Re}\lambda_j|^{1/3}\} < C. \tag{4}$$

This is known to be optimal in the following sense [2]: a non-degenerate isolated simple geodesic γ of length d_γ on the boundary of an analytic strictly convex obstacle in $\mathbb{R}^{n+1}$, $n+1$ odd, generates infinitely many poles in any region

$$\{z:\ \operatorname{Im} z < \frac{B}{d_\gamma}|z|^{\frac{1}{3}}\},\ B > \omega_\gamma = 2^{-\frac{1}{3}}\zeta_1 \cos\left(\frac{\pi}{6}\right) \cdot \int_0^{d_\gamma} \rho^{2/3}(s)ds,$$

where $-\zeta_1$ is the first zero of the Airy function, ρ is the curvature of γ in $\mathbb{R}^{n+1}$ and s the arc length parameter.

For strictly convex obstacles the results on pole counting are expected to be more precise and in particular the density of poles near the real axis has already been estimated. To motivate them let us consider (in [28] the convention was that $\operatorname{Im}\lambda_j < 0$)

$$N_\theta(r) := \sharp\{\lambda_j:\ \lambda_j \text{ is a pole of } R_{\mathcal{O}}(\lambda),\ |\lambda_j| \leq r,\ 0 \leq \arg\lambda_j < \theta,\ \pi - \theta < \arg\lambda_j \leq \pi + \theta\}.$$

Then for strictly convex smooth obstacles we get

$$N_\theta(r) = \mathcal{O}(\theta^{3/2})r^{n+1},\ r > 1. \tag{5}$$

By applying complex scaling "all the way to the boundary" which is possible because of strict convexity, but still using a form of functional calculus from [20], (5) was obtained in [21] for an arbitrary convex obstacle provided $r > r(\theta)$. It was noted by Hargé and Lebeau [7] that a particular choice of the angle of scaling yields, among other things, a cubic pole free region of the form (4) for strictly convex obstacles with smooth boundaries. It had been widely believed in applied mathematics and known already in dimensions two and three (see [1]).

1.2. **Trapping obstacles: results of Ikawa and Gerard.**

We consider scattering by two strictly convex obstacles. Note that scatterer is always trapping if it is not connected.

We suppose that

$$\begin{aligned} &\mathcal{O} = \Omega_1 \cup \Omega_2 \subset \mathbb{R}^{n+1},\ \text{with } C^\infty \text{ boundary } \partial\Omega_i,\ i = 1, 2, \\ &\text{where } \Omega_1,\ \Omega_2 \text{ are compact and } \textit{strictly convex},\ \Omega_1 \cap \Omega_2 = \emptyset. \end{aligned} \tag{6}$$

Let d be the distance between Ω_1 and Ω_2 and $a_i \in \partial\Omega_i$ the points on the boundary such that $|a_1 - a_2| = d$. Then there is one trapped broken characteristic ray connecting a_1 and a_2.

Under these assumptions on Ω Bardos, Guillot and Ralston (*cf.* [3], for $n+1$ odd) show the existence of an infinite number of resonances in $\{z;\ \operatorname{Im} z \leq \epsilon \log|z|\}$ for any $\epsilon > 0$.

Thus their result shows a difference in location of resonances between cases of trapping obstacles and of non-trapping obstacles.

The most complete results on location of poles were given by Ikawa and Gérard. There results shows that uniformly in the strip $0 < \operatorname{Im} z < c$, as $\operatorname{Re} z$ goes to infinity the resonances are well approximated by pseudo-poles of [3]:

$$\lambda_{\alpha,k} = k\frac{\pi}{d} + \frac{i}{4d}\sum_{j=1}^{n} \ln\nu_j \cdot (2\alpha_j+1),\ (k,\alpha)\in\mathbb{Z}\times\mathbb{N}^n, \tag{7}$$

where ν_j are the eigenvalues of the linear Poincaré map $\mathrm{D}\kappa(a_1,0)$.

Though Ikawa and Gérard only consider the case of odd dimensions $n+1$, their results are valid in even dimensions $n+1\geq 2$ as well.

In [15] Ikawa obtained the first string of resonances closest to the real axis ($\alpha=0$) and in (*cf.* [9]) Gérard got the complete asymptotics of all strings.

We start by defining the canonical transformation of the billiard $\kappa:\ T^*\partial\Omega_1\mapsto T^*\partial\Omega_1$ following Gérard [9]. For $\rho\in T^*\partial\Omega_1$ in a neighborhood of $(a_1,0)\in T^*\partial\Omega_1$ we draw an outgoing half-ray γ issued from ρ and we denote γ' the half-ray reflected from $\partial\Omega_2$. If γ' intersects $\partial\Omega_1$ then we define $\kappa(\rho)$ the projection on $T^*\partial\Omega_1$ of the point of intersection between γ' and $\partial T^*(\mathbb{R}^{n+1}\setminus\Omega_1)$.

For $\rho\in T^*\partial\Omega_1$, close to $(a_1,0)$, we denote by $\kappa_1(\rho)$ the projection on $T^*\partial\Omega_2$ of the point of intersection of γ and $\partial T^*(\mathbb{R}^{n+1}\setminus\Omega_2)$. κ_1 is the canonical transformation from $T^*\partial\Omega_1$ to $T^*\partial\Omega_2$. If we define in the similar way the canonical transformation $\kappa_2:\ T^*\partial\Omega_2\mapsto T^*\partial\Omega_1$ then the billiard map is the canonical transformation

$$\kappa=\kappa_2\circ\kappa_1. \tag{8}$$

According to Petkov ([17]), [18], Corollary 2.3.3) and Bardos, Guillot, Ralston ([3], Proposition 3) using that Ω_1, Ω_2 are strictly convex, the eigenvalues of $\mathrm{D}\kappa(\rho_1,0)$ are positive $\neq 1$, thus κ is of hyperbolic type and we have $\kappa(a_1,0)=(a_1,0)$.

Denote by ν_j the eigenvalues >1 of $\mathrm{D}\kappa(a_1,0)$ numerated such that $1<\nu_1\leq\nu_2\leq\ldots\leq\nu_n$ and put $b_0=\Pi_{i=1}^{n}\nu_j^{-1/2}$. For $\alpha\in\mathbb{N}$ we put $K_\alpha=b_0\nu^{-\alpha}$. (It is possible that $K_\alpha=K_{\alpha'}$ for $\alpha\neq\alpha'$). For any value of K_α and $a_{\alpha,j,l}\in\mathbb{C}$, we define

$$\lambda_l(\alpha,k)=\lambda_{\alpha,k}+\sum_{j=1}^{\infty}a_{\alpha,j,l}\lambda_{\alpha,k}^{-j/2al}\ \text{with}\ al\in\mathbb{N},\ l=1,\ldots,a \tag{9}$$

which correspond to the asymptotic expansion for the eigenvalues of an $a\times a$ matrix, where

$$a=\mathrm{Card}\,\{\alpha'|\ K_{\alpha'}=K_\alpha\}. \tag{10}$$

Let p_l be the multiplicity of $\lambda_l(\alpha,k)$ as asymptotic eigenvalue.

Theorem 1 (Gérard, $n+1$ odd)**.** *For all $A>0$, there exists $C>0$ such that, if $\lambda_{\alpha,k}$ is given by (7), then there exist coefficients $a_{\alpha,j,l}\in\mathbb{C}$ in asymptotic expansions (9) such that for all $N\in\mathbb{N}$ there exist $k_N\in\mathbb{N}$ and $c_N\in\mathbb{R}$ such that the poles situated in the region*

$$\{\lambda\in\mathbb{C};\ \mathrm{Im}\,\lambda\leq A,\ |\lambda|>C\}$$

are all in the balls

$$\left|\lambda-\left(\lambda_{\alpha,k}+\sum_{j=1}^{m_N}a_{\alpha,j,l}\lambda_{\alpha,k}^{-j/2al}\right)\right|\leq C_N|\lambda_{\alpha,k}|^{-N},\ k\geq k_N\in\mathbb{N},$$

where m_N is the largest j such that $\frac{j}{2al}<N$, a is defined in (10) and in each ball there are exactly p_l poles with multiplicities.

In his article [9] Gérard reduces the problem to the problem on the boundary of the one obstacle, $\partial\Omega_1$, by introducing the quantum billiard operator $M(\lambda)$ which quantizes the non-linear Poincaré transform along the trapped trajectory κ. The operator $M(\lambda)$ is defined in the following way.

Let $H_{i,+}(\lambda):\ C^\infty(\partial\Omega_i)\mapsto C^\infty(\mathbb{R}^{n+1}\setminus\Omega_i),\ i=1,2$, be the outgoing resolvent of the problem

$$\left\{\begin{array}{l}(\Delta+\lambda^2)H_{i,+}(\lambda)v=0 \text{ in } \mathbb{R}^{n+1}\setminus\Omega_i\\ H_{i,+}(\lambda)v_{|\partial\Omega_i}=v\end{array}\right. \tag{11}$$

extended as an operator $H^{1/2}(\partial\Omega_i)\mapsto H^1_{\rm loc}(\complement\Omega_i)$ and analytical for $\operatorname{Im}\lambda<0$. It is known that $H_{i,+}(\lambda)$ extends analytically as a bounded operator $H^{1/2}(\partial\Omega_i)\mapsto H^1_{\rm loc}(\complement\Omega_i)$ to a domain of the form (1) if the boundary $\partial\mathcal{O}$ is C^∞ and (3) if the boundary is real analytic.

We define $H_i(\lambda)v=H_{i,+}(\lambda)v_{|\partial\Omega_{i+1}}$, where $\partial\Omega_3=\partial\Omega_1$ and

$$M(\lambda)=H_2(\lambda)H_1(\lambda)=\gamma_1H_{2,+}\gamma_2H_{1,+}:\ H^1(\partial\Omega_1)\mapsto H^1(\partial\Omega_1), \tag{12}$$

where $\gamma_i:\ H^1_{\rm loc}(\complement\Omega_i)\mapsto H^{1/2}(\partial\Omega_i)$ is the operator of restriction to $\partial\Omega_i$.

The operator $M(\lambda)$ defined on $H^1(\partial\Omega_1)$ for $\operatorname{Im}\lambda<0$ continues analytically as a bounded operator $H^{1/2}(\partial\Omega_1)\mapsto H^{1/2}(\partial\Omega_1)$ in the domain of the form (1) if $\partial\mathcal{O}$ is C^∞ and (3) if $\partial\mathcal{O}$ is real analytic, and there satisfies the following estimate

$$\exists D>0,\ \exists C>0,\ \|M(\lambda)\|_{\mathcal{L}(H^{1/2}(\partial\Omega_1))}\le C|\lambda|^2e^{D\operatorname{Im}\lambda^+},$$

where $\operatorname{Im}\lambda^+=\max(\operatorname{Im}\lambda,0)$.

We have the following relation between the outgoing resolvent $H_+(\lambda)$ in the exterior of $\mathcal{O}=\Omega_1\cup\Omega_2$ ($u=H_+v$ satisfies (11) with $\partial\Omega$ instead of $\partial\Omega_i$) and the billiard operator M, used by Gérard in [9], page 91:
if $(v_1,v_2)\in\ C^\infty(\partial\Omega_1)\times C^\infty(\partial\Omega_2)$ then

$$H_+(\lambda)(v_1,v_2)=(H_{1,+}-H_{2,+}H_1)(I-M)^{-1}v_1+(H_{2,+}-H_{1,+}(I-M)^{-1}H_2+H_{2,+}H_1(I-M)^{-1}H_2)v_2.$$

Using this relation Gérard proved in [9] that

Lemma 1. *The scattering poles counted with their multiplicities coincide with*

$$\{\lambda,\ 0\in\sigma(I-M(\lambda))\}. \tag{13}$$

Strictly speaking, Gérard only considers the case of λ in a strip

$$0\le\operatorname{Im}\lambda\le c_1,\ \operatorname{Re}\lambda\ge c_2, \tag{14}$$

for c_2 sufficiently large and $n+1$ odd, but his proof also works in more general domains (1), (3).

Microlocally near a_1, Gérard reduced the problem (13) to the problem of finding the points λ in a strip (14) for which the operator $I-e^{-2id\lambda}M_0(\lambda)$ considered in some appropriate space has non-trivial kernel. Here $M_0(\lambda)$ is a semi-classical Fourier integral operator associated to the real hyperbolic canonical transformation κ with a fixed point at $(0,0)$. Gérard obtained his result by approximating of $M_0(\lambda)$ by its linearization.

In the case when $\nu_j,\ j=1,\dots,n$ are linearly independent over the field of integer numbers and the boundaries are analytic, we will use the full Birkhoff normal form of M_0 (as in [13]) in order to get the explicit formulas for the resonances in any logarithmic neighborhood of the real axis of the form (1).

2. OUR RESULT.

We suppose in addition to (6) that the boundaries $\partial\Omega_i,\ i=1,2$, are real analytic. Suppose that the eigenvalues of the Poincaré map

$$0<\nu_n^{-1}\le\ldots\le\nu_1^{-1}<1<\nu_1\le\ldots\le\nu_n$$

verify the non-resonance condition:

$$\sum_{1}^{n} k_j \ln \nu_j = 0,\ k_j \in \mathbb{Z} \implies k_1 = \ldots = k_n = 0. \tag{15}$$

In order to have good separation of the strings of resonances we impose the Diophantine condition:

$$\alpha \neq \beta,\ |\alpha|, |\beta| \leq m \ \Rightarrow\ |\ln \nu(\alpha - \beta)| \geq \frac{1}{C(D)} e^{-Dm},\ D > 0. \tag{16}$$

Then microlocally near $a_1 = 0$ we can transform $M_0(\lambda)$ into its semiclassical Birkhoff normal form to any order r

$$M_{00}(\lambda) := e^{-i\lambda F^r(I_1,\ldots,I_n;1/\lambda)},\ I_j = \left(x_j \partial_{x_j} + 1/2\right)/i\lambda.$$

Here for any $r \in \mathbb{Z}_+$, and $\imath$ in a neighborhood of $\imath = 0$ in $\mathbb{R}^n$,

$$F^r(\imath; 1/\lambda) = F_0(\imath) + \frac{1}{\lambda} F_1(\imath) + \frac{1}{\lambda^2} F_2(\imath) + \ldots + \frac{1}{\lambda^r} F_r(\imath) \tag{17}$$

with F_j polynomial of degree $(r - j)$ in $\imath$, with

$$F_0(\imath) = G(\imath, \mu) + H(\imath),\ G(\imath, \mu) = \sum_{i=1}^{n} \mu_i \imath_i,\ \mu_i = \ln \nu_i,\ H(\imath) = \sum_{j=2}^{r} h_j(\imath) = O(\imath^2),$$

$$\text{where } h_j(\imath) = \sum_{|\alpha|=j} \frac{1}{\alpha!} \partial^\alpha H(0) \imath^\alpha \text{ is a homogeneous polynomial of degree } j.$$

Here $F_0(\imath)$ is real when $\imath \in \mathbb{R}^n$.

We have for any $r, r' \geq j$:

$$F_j^r(\imath) - F_j^{r'}(\imath) = \mathcal{O}\left(\imath^{\min\{r,r'\}-j+1}\right).$$

In other words, $h_j(\imath)$ do not depend on r.

Thus F^r, F_j are holomorphic in a neighborhood of $\imath = 0$ and $M_{00}(\lambda)$ is analytic in any logarithmic neighborhood of the real axis of the form

$$\Lambda_{A,B} := \{\lambda \in \mathbb{C} \text{ or } \Lambda;\ \operatorname{Im} \lambda < A \ln \operatorname{Re} \lambda,\ \operatorname{Re} \lambda > B\}. \tag{18}$$

Let

$$P(\lambda) := I - e^{-2id\lambda} M_{00}(\lambda).$$

As $I_j x_j^{\alpha_j} = \dfrac{1}{2i\lambda}(2\alpha_j + 1) x_j^{\alpha_j}$, the monomials $x^\alpha = x_1^{\alpha_1} \cdot x_2^{\alpha_2} \cdot \ldots \cdot x_n^{\alpha_n}$ are formally the eigenfunctions of the operator $P(\lambda)$. We have then

$$P(\lambda) x^\alpha = \left(1 - e^{-2id\lambda} K_\alpha(\lambda)\right) x^\alpha,\ K_\alpha(\lambda) := e^{-i\lambda F^r\left(\frac{2\alpha+1}{2i\lambda}; 1/\lambda\right)}.$$

We approximate the problem (13) with the following model problem:

$$1 - e^{-2id\lambda} K_\alpha(\lambda) = 0 \ \Leftrightarrow\ 2d\lambda + \lambda F^r\left(\frac{2\alpha + 1}{2i\lambda}; 1/\lambda\right) = 2\pi k,\ \alpha \in \mathbb{Z}_+^n,\ k \in \mathbb{Z},\ \lambda \in \Lambda_{A,B}. \tag{19}$$

Note that by substituting F^r by its leading term near the origin, $\sum_{i=1}^n \ln \nu_i \cdot \imath_i$, we recover the pseudopoles (7).

In general in order to solve (19) we observe that if $F(\imath;1/\lambda)=F_0(\imath)+F_1(\imath)/\lambda+F_2(\imath)/\lambda^2+\dots$ is either an asymptotic or a finite sum, and F, F_j are holomorphic in a fixed neighborhood of $\imath=0$, and $F_0(0)$ is small (in our case $F_0(0)=0$), then

$$\frac{\partial}{\partial\lambda}\left(2d\lambda+\lambda F\left(\frac{2\alpha+1}{2i\lambda};1/\lambda\right)\right)=$$
$$=2d+F\left(\frac{2\alpha+1}{2i\lambda};1/\lambda\right)-\frac{1}{\lambda}\left(\frac{\partial F\left(\frac{2\alpha+1}{2i\lambda};1/\lambda\right)}{\partial\imath}\frac{2\alpha+1}{2i}-\frac{\partial F\left(\frac{2\alpha+1}{2i\lambda};1/\lambda\right)}{\partial h}\right)\approx 2d$$

for $|\alpha|/|\lambda|$ small.

We omit r and write equation (19) in the form

$$\frac{\lambda}{k}\left(1+\frac{1}{2d}F\left(\frac{(2\alpha+1)/k}{2i\lambda/k};\left(\frac{\lambda}{k}\right)^{-1}\frac{1}{k}\right)\right)=\frac{\pi}{d} \tag{20}$$

with λ/k as unknown variable.

It is then clear that equation (20) has the solution

$$\frac{\lambda_0}{k}=g(\frac{\alpha}{k};\frac{1}{k}),\ g(\theta;\frac{1}{k})=\frac{\pi}{d}+g_0(\theta)+g_1(\theta)\frac{1}{k}+\dots,\ g_0(0)=0.$$

We get the solution in the form

$$\frac{\lambda_0}{k}=\frac{\pi}{d}+\frac{a_1}{k}+\frac{a_2}{k^2}+\dots=\frac{\pi}{d}-\frac{F_1(0)}{2dk}+\frac{\mathrm{Re}\,(a_2)}{k^2}+\dots+i\left(\frac{1}{4dk}\sum_{j=1}^{n}\mu_j(2\alpha_j+1)+\frac{\mathrm{Im}\,(a_2)}{k^2}+\dots\right),$$

for $k>>1$, $a_j=O(|\alpha|^j)$, $\alpha=\mathcal{O}(\ln k)$, where a_j for $j=1,2,\dots$ are polynomials in α (see Theorem 5 for details) and $F_1(0)$ is real constant as the operator M is microlocally unitary (see Appendix A.2). The coefficient a_j is only dependent on the coefficients in the Birkhoff normal form of order j, $F^j(\imath;1/\lambda)=F^j_0(\imath)+\frac{1}{\lambda}F^j_1(\imath)+\frac{1}{\lambda^2}F^j_2(\imath)+\dots+\frac{1}{\lambda^r}F^j_r(\imath)$, and independent of r for $j\le r$. If λ_0^r and $\lambda_0^{r'}$ are two such solutions with $r<r'$ then

$$\frac{\lambda_0^{r'}}{k}-\frac{\lambda_0^{r}}{k}=\frac{a^{r'}_{r+1}-a^{r}_{r+1}}{k^{r+1}}+\mathcal{O}\left(\frac{|\alpha|^{r+2}}{k^{r+2}}\right)=\mathcal{O}\left(\frac{|\alpha|^{r+1}}{k^{r+1}}\right)$$

The main result of the paper is the following theorem:

Theorem 2. *Suppose (15) and (16) are satisfied and let $\Lambda_{A,B}$ be the logarithmic neighborhood of real axis as in (18). For any $N\in\mathbb{N}$ and for all $A>0$ there exists $B>0$, and $r=r(N)\in\mathbb{N}$ such that there exist functions $a_j=a_j(\alpha)$, $j=1,\dots,r$, polynomial in α of order $\le j$ ($a_j=\mathcal{O}\left(|\alpha|^j\right)$ for $|\alpha|\ge 1$) and there exists bijection b_N between the set of*

$$\lambda^r_{\alpha,k}=k\left(\frac{\pi}{d}+\frac{a_1}{k}+\frac{a_2}{k^2}+\dots+\frac{a_{r+1}}{k^{r+1}}\right)\in\Lambda_{A,B}$$

and the set of resonances in $\Lambda_{A,B}$, where elements in each set are counted with their multiplicity, such that

$$b_N(\lambda)-\lambda=\mathcal{O}\left(|\lambda|^{-N}\right).$$

Solutions to (13) can be given by means of the Grushin problem for $I-M$ in some weighted Sjöstrand's space $H(\Lambda_{tG})$ associated to the escape function G. The results of propagation of analytic singularities by Lebeau [16] can be applied to localize the analytic singular support of M. Outside the analytic singular support the norm of M is small. In a neighborhood of the analytic singular support we use the properties of the escape function. Outside some small neighborhood V_0 of $a_1\in\partial\Omega_1$ we get that the operator $I-M(\lambda)$ is invertible. In V_0 we reduce M

by means of an analytic Fourier integral operator of Bargman type U to the operator $e^{-2id\lambda}M_0(\lambda)$ in the Birkhoff normal form such that for any $N_0 \in \mathbb{N}$,

$$UM(\lambda)u = e^{-2id\lambda}M_0(\lambda)Uu + \mathcal{O}\left(|\lambda|^{-N_0}\right)|u|_{\mathcal{H}} \text{ in } H_\Phi.$$

Here $M_0 \equiv e^{-i\lambda F(I;1/\lambda)}$ in the sense of [13]. We use the truncated Birkhoff normal form to the order r, $e^{-i\lambda F^r(I;1/\lambda)}$, which is an analytic Fourier integral operator and the Grushin problem in order to show that the solutions of (13) can be approximated by the solutions of the model problem

$$\{\lambda \in \mathbb{C},\ 0 \in \sigma\left(I - e^{-2id\lambda}e^{-i\lambda F^r(I;1/\lambda)}\right)\} \tag{21}$$

which leads to (19).

Note that as $R_{\mathcal{O}}(-\overline{\lambda}) = \overline{R_{\mathcal{O}}(\lambda)}$ it is sufficient to consider the case $\operatorname{Re}\lambda > 0$.

The structure of the paper is the following:

In Section 3 we construct the escape function.

In Section 4 we review the method of FBI-Bargman transforms.

In Section 5 we apply the FBI-Bargman transform to the quantum billiard operator M and consider the norm of the transformed operator M_1.

In Section 6 we reduce M_1 to the Birkhoff normal form up to some order r at a_1.

In Section 7 we deform the space near a_1 such that monomials form an almost orthonormal basis in some neighborhoodof a_1.

In Section 8 we derive the asymptotic expansions for the solutions of the model problem (13).

In the last sections 9-12 we prove Theorem 2 by the routine method of Grushin problems.

In Appendix A we review the norm estimates of the quantum billiard operator M in the Sobolev space and show that M is microlocally unitary with respect to the flux norm.

Acknowledgements. The author is grateful to Jöhannes Sjöstrand for numerous discussions and constant support during the preparation of the manuscript.

3. Construction of the global escape function on $X = T^*\partial\Omega_1$ for the billiard canonical transformation κ.

3.1. **Definitions of the domains.**

Let $\kappa:\ T^*\partial\Omega_1 \mapsto T^*\partial\Omega_1$ be the billiard map defined in (8). We have $\kappa(a_1,0) = (a_1,0)$. For $\rho = (x,\xi) \in \operatorname{neigh}(a_1,0) \subset T^*\partial\Omega_1$ we draw an outgoing half-ray γ issued from ρ. If γ intersects $\partial T^*(\mathbb{R}^{n+1}\setminus\Omega_2)$ then we denote γ' the reflected half-ray. If γ' intersects $\partial T^*(\mathbb{R}^{n+1}\setminus\Omega_1)$ then we define $\kappa(\rho)$ the projection on $T^*\partial\Omega_1$ of the point of intersection between γ' and $\partial T^*(\mathbb{R}^{n+1}\setminus\Omega_1)$. Let $D(\kappa) = \{\rho \in T^*\partial\Omega_1;\ \exists\kappa(\rho) \in T^*\partial\Omega_1\}$ be the domain of definition of κ.

We denote $X = \partial\Omega_1$. The points in T^*X can be divided into the following 3 regions:

$$\begin{aligned}&\mathcal{H} \text{ hyperbolic region: } \{(y,\eta) \in T^*X;\ |\eta| < 1\}\\ &\mathcal{G} \text{ glancing region: } \{(y,\eta) \in T^*X;\ |\eta| = 1\}\\ &\mathcal{E} \text{ elliptic region: } \{(y,\eta) \in T^*X;\ |\eta| > 1\}.\end{aligned}$$

We denote $\rho_1 = (a_1,0) \in T^*X$ and $B(\rho_1,c) := \{\rho \in T^*X;\ \operatorname{dist}(\rho,\rho_1) < c\}$.

We use the following convention: h denotes $1/\operatorname{Re}\lambda$, $\lambda \in \Lambda_{A,B}$.

Let $W_0 \subset T^*X$ be a small neighborhood of ρ_1 of the size $\sqrt{h\ln(1/h)}$ given by

$$W_0 := B(\rho_1, c_0\sqrt{h\ln(1/h)}) \subset T^*\partial\Omega_1, \tag{22}$$

for some constant $c_0 > 0$.

Let $W_1 \subset T^*X$ be a larger neighborhood of ρ_1 independent of h such that

$$W_0 \subset\subset W_1 \subset B(\rho_1, c_1) \subset D(\kappa) \subset \mathcal{H}$$

for some constant $c_1 > 0$ (it will be defined later in (26)).

Proposition 1. *There exists a real function $G \in C^\infty(T^*\partial\Omega_1)$ with the following properties:*

1. $G = \mathcal{O}\left(h\ln^2(1/h)\right)),\ |\nabla G| = \mathcal{O}\left(\sqrt{h\ln(1/h)}\right),\ |\nabla^2 G| = \mathcal{O}(1)$;
2. *for all* $\rho \in W_0,\ G(\kappa(\rho)) - G(\rho) \geq C\,\mathrm{dist}(\rho_1, \rho)^2$;
3. *for all* $\rho \in W_1 \setminus W_0,\ G(\kappa(\rho)) - G(\rho) \geq C\,h\ln(1/h)$;
4. *for* $\rho \in T^*\partial\Omega_1 \setminus W_1,\ G(\kappa(\rho)) - G(\rho) \geq C\,h\ln^2(1/h)$.

Here C is some positive constant.

We call G a (global) escape function for κ. In the following sections we prove Proposition 1.

3.2. **Construction of the local escape function G_{int}.** As κ is of hyperbolic type, by the stable manifold theorem, in some neighborhood W of $(a_1, 0)$, κ has an incoming stable manifold Λ_- and an outgoing stable manifold Λ_+ which are lagrangian manifolds, intersecting transversally at $(a_1, 0)$ characterized by

$$\Lambda_\pm = \{(x,\xi) \in W|\ \kappa^{-n}(x,\xi) \in W \text{ for all } n \in \mathbb{Z}_\pm\}$$

and if $(x,\xi) \in \Lambda_\pm$ then $\kappa^{-n}(x,\xi) \mapsto (a_1, 0)$ exponentially fast as $n \to \pm\infty$.



FIGURE 1. *A part of $T^*\partial\Omega_1$ near $(a_1, 0)$.*

There exists a local canonical transformation $f : \mathrm{neigh}((a_1,0), T^*X) \mapsto \mathrm{neigh}(0, \mathbb{R}^{2n})$ such that $f(\Lambda_+) = \{\xi = 0\}$, $f(\Lambda_-) = \{x = 0\}$ in a neighborhood W of $(a_1, 0)$ which corresponds to the symplectic change of coordinates. In these new coordinates the differential of κ at $(0,0)$ has the form

$$\mathrm{D}\kappa(0,0) = \begin{pmatrix} A & 0 \\ 0 & (A^T)^{-1} \end{pmatrix},$$

the eigenvalues of A are $\nu_1, \ldots, \nu_n$.

By the theorem of Lewis and Sternberg (*cf.* [27], [6]) there exists a Hamiltonian $p(x,\xi)$ such that $\kappa \equiv \exp H_p$ in the sense of formal Taylor series (*cf.* [12] [13]) and p has the form $p(x,\xi) = B(x,\xi)x\cdot\xi$, where $B(0,0) = \ln A$ with eigenvalues $\mu_j = \ln \nu_j$, $j = 1, \ldots, n$.

When κ is of the form $\exp H_p$ near $(0,0)$ and $G \in C^\infty$ with $dG(0,0) = 0$, then locally in a neighborhood of $(0,0)$ we get by Taylor expansion

$$G(\exp H_p(\rho)) = \sum \frac{1}{k!} H_p^k G(\rho) = G(\rho) + H_p G + \mathcal{O}\left((x,\xi)^3\right)$$

and $G(\kappa(\rho))-G(\rho)=H_pG+\mathcal{O}\left((x,\xi)^3\right)$. Thus G with properties as in Proposition 1 is an escape function for p in the usual sense as defined in [10]. This can be used in order to construct G locally near 0.

Suppose that the coordinates are chosen such that

$$\kappa=\exp H_p,\ p=\sum_{i=1}^{n}\mu_i x_i\xi_i+\mathcal{O}\left((x,\xi)^3\right).$$

It is enough to construct an escape function for $p:=x\xi,\ x,\xi\in\mathbb{R}$.

Inside the ball $x^2+\xi^2<\mathcal{O}(1)h\ln(1/h)$ we can take

$$G_0(x,\xi)=\frac{1}{2}(x^2-\xi^2).$$

Then, $H_pG_0=x^2+\xi^2=\mathcal{O}\left(h\ln(1/h)\right),\ \nabla G_0=\mathcal{O}\left(\sqrt{h\ln(1/h)}\right),\ G_0=\mathcal{O}\left(h\ln(1/h)\right)$.

For $c_1h\ln(1/h)\leq x^2+\xi^2\leq c_2$ we use the following ansatz: $G_1=f_h(x)-f_h(\xi)$.

We have $H_p=x\partial_x-\xi\partial_\xi$ and we want that $H_pG_1\sim h\ln(1/h)$ in this region.

We can for example choose $f_h(x)$ satisfying

$$\begin{cases} x\partial_x f_h(x)=h\ln(1/h), & \sqrt{h\ln(1/h)}<|x|<c\\ f_h(\sqrt{h\ln(1/h)})=h\ln(1/h)/2 \end{cases}$$

which gives $f_h(x)=h\ln(1/h)\ln\left(\frac{x}{\sqrt{h\ln(1/h)}}\right)+\frac{h\ln(1/h)}{2}$. And similarly for $f_h(\xi)$.

With the above example in mind we define a local escape function in the following way.

Let f be a local real canonical transformation $f(y,\eta)=(x,\xi)$ such that Λ_- is transformed into $\{x=0\}$ and $\Lambda_+\mapsto\{\xi=0\}$ and let $\widehat{W_1}$ be the image of W_1,

$$f(W_1)=\widehat{W_1}\subset\operatorname{neigh}(0,\mathbb{R}^{2n}).$$

Put $s=h\ln(1/h)$.

The local escape function G_1 on $\widehat{W_1}$ is defined by $G_1=G_+-G_-$, where

$$G_+=\frac{1}{2}s\ln\left(s+x^2\right),\ G_-=\frac{1}{2}s\ln\left(s+\xi^2\right).$$

Then

$$G_1=\frac{1}{2}s\ln\left(\frac{s+x^2}{s+\xi^2}\right).$$

If $x^2<<s$ then

$$G_+=\frac{1}{2}s\left\{\ln(s)+\ln\left(1+\frac{x^2}{s}\right)\right\}=\frac{1}{2}s\ln(s)+\frac{1}{2}s\cdot\frac{x^2}{s}+s\mathcal{O}\left(\left(\frac{x^2}{s}\right)^2\right).$$

We have then for $x^2+\xi^2<<s$,

$$G_1=G_+-G_-=\frac{1}{2}s\ln\left(\frac{s+x^2}{s+\xi^2}\right)=\frac{1}{2}(x^2-\xi^2)+\mathcal{O}\left(\frac{(x^2+\xi^2)^2}{s}\right). \tag{23}$$

If $s<<x^2\leq C$, $\xi=0$, we get

$$G_1=\frac{1}{2}s\ln\left(\frac{x^2(1+\frac{s}{x^2})}{s}\right)=\frac{1}{2}s\ln\left(\frac{x^2}{s}\right)+\mathcal{O}\left(\frac{s^2}{x^2}\right)=\frac{1}{2}h(\ln(1/h))^2+\mathcal{O}\left(h\ln(1/h)\ln\ln(1/h)\right).$$

Thus the variation of G_1 is of order $h\ln^2(1/h)$.

For the derivatives we get

$$\partial_x G_1(x,\xi) = \frac{sx}{s+x^2},\ \partial_\xi G_1(x,\xi) = -\frac{s\xi}{s+\xi^2}.$$

As

$$\partial_x^2 G_1(x,\xi) = \frac{s}{s+x^2}\left(1-\frac{2x^2}{s+x^2}\right)$$

the maximum of $\partial_x G_1$ is attained when $x^2 = s$ with the maximum value $\sqrt{s}/2$. Using the similar estimate for $\partial_\xi G_1$, we get

$$|\nabla G_1| = \mathcal{O}\left(\sqrt{s}\right).$$

We have

$$\partial_x^3 G_1(x,\xi) = -\frac{2sx}{(s+x^2)^2}\left(3-\frac{4x^2}{s+x^2}\right)$$

and we get that $\partial_x^2 G_1$ attains the minimum at $x^2 = 3s$ with the minimum value $-1/8$.

By the similar estimate for $\partial_\xi^2 G_1$, we get

$$|\nabla^2 G_1| = \mathcal{O}(1).$$

Similarly we get $|x\partial_x G_1| = \mathcal{O}(s)$, $|\xi\partial_\xi G_1| = \mathcal{O}(s)$.

We denote the billiard canonical transformation in the new coordinates by $\hat{\kappa} := f\circ\kappa\circ f^{-1} : \widehat{W}_1 \mapsto \widehat{W}_1$. Denote $\rho = (x,\xi)$. We have proved the first statement in

Lemma 2. *There exists a real function $G_1 \in C^\infty(\widehat{W}_1)$ with the following properties:*

1. $G_1 = \mathcal{O}\left(h\ln^2(1/h)\right)$, $|\nabla G_1| = \mathcal{O}\left(\sqrt{h\ln(1/h)}\right)$, $|\nabla^2 G_1| = \mathcal{O}(1)$;
2. *for all* $\rho\in\widehat{W}_1$, $|\rho| \le c\sqrt{h\ln(1/h)}$, $G_1(\hat{\kappa}(\rho)) - G_1(\rho) \ge C\,|\rho|^2$;
3. *for all* $\rho\in\widehat{W}_1$, $|\rho| \ge c\sqrt{h\ln(1/h)}$, $G_1(\hat{\kappa}(\rho)) - G_1(\rho) \ge C\,h\ln(1/h)$.

Here c, C denote different positive constants.

Proof of (2) and (3): The canonical transformation in the new coordinates is of the form $\hat{\kappa}(x,\xi) = (Ax, (A^T)^{-1}\xi) + \mathcal{O}((x,\xi)^2)$, where A is diagonal with entries e^{μ_i}, $i = 1,\dots,n$.

We put $n = 1$, extension to the general case is straightforward.

Let $s := h\ln(1/h)$. Then, choosing $\widehat{W}_1$ small enough, we have for $\rho\in\widehat{W}_1$,

$$G_1(\rho) = \frac{1}{2}s\ln\left(\frac{s+x^2}{s+\xi^2}\right),\ G_1(\hat{\kappa}(\rho)) = \frac{1}{2}s\ln\left(\frac{s+e^{2\mu}x^2}{s+e^{-2\mu}\xi^2}\right) + \mathcal{O}\left(s(x,\xi)\right).$$

If $|\rho|^2 = x^2+\xi^2 \le cs$, then we can use the asymptotics (23):

$$G_1(\hat{\kappa}(\rho)) - G_1(\rho) = (e^{2\mu}-1)x^2 - (e^{-2\mu}-1)\xi^2 + \mathcal{O}\left(\frac{|\rho|^4}{s}\right) \ge 2\mu|\rho|^2 + \mathcal{O}\left(\frac{|\rho|^4}{s}\right).$$

If $cs \le |\rho|^2 = x^2+\xi^2 \le \epsilon$, for ϵ small enough, we have

$$G_1(\hat{\kappa}(\rho)) - G_1(\rho) = \frac{1}{2}s\left(\ln\left(\frac{s+e^{2\mu}x^2}{s+x^2}\right) + \ln\left(\frac{s+\xi^2}{s+e^{-2\mu}\xi^2}\right)\right) + \mathcal{O}\left(s|\rho|\right) \ge \tag{24}$$

$$\ge \frac{1}{2}s\left(\ln\left(\frac{1+e^{2\mu}c}{1+c}\right) + \ln\left(\frac{1+c}{1+e^{-2\mu}c}\right)\right) + \mathcal{O}\left(s|\rho|\right) \ge C_\epsilon s,$$

using that the right hand side of (24) is increasing function in x and in ξ. Here $C_\epsilon > 0$ depends on the size of order ϵ of $\widehat{W}_1$. ∎

In the original space (in coordinates (y,η)) in a neighborhood $W_1 \subset T^*\partial\Omega_1$ we put

$$G_{\mathrm{int}}(y,\eta) = G_1(x,\xi),\ f(y,\eta) = (x,\xi) \ \Leftrightarrow\ G_{\mathrm{int}} := G_1\circ f^{-1}.$$

We call $G_{\rm int}$ the local escape function for κ.

3.3. Construction of the exterior escape function $G_{\rm ext}$.

As before $D(\kappa)$ denotes the domain of definition of κ and

$$R(\kappa) := \{\kappa(\rho),\ \rho \in D(\rho)\}$$

the image of κ. On $R(\kappa)$ we can define the inverse κ^{-1}.

We define

$$\widetilde{\Omega}_+(0) = \overline{\mathcal{H}} = \mathcal{H} \cup \mathcal{G},\ \widetilde{\Omega}_+(1) = D(\kappa),\ \text{for } j \geq 1,\ \widetilde{\Omega}_+(j) := D(\kappa^j),$$
$$\widetilde{\Omega}_+(0) \supset \widetilde{\Omega}_+(1) \supset \widetilde{\Omega}_+(2) \supset \dots,\ \bigcap_{j=0}^{\infty} \widetilde{\Omega}_+(j) = \Lambda_-^{\rm ext},$$
$$\Lambda_-^{\rm ext} = \{(x,\xi) \in T^*X,\ \forall j \in \mathbb{N},\ \exists \kappa^j(x,\xi) \in T^*X\},$$

where ext stands for "extended" by the iterated action of κ. Note that as the obstacles are strictly convex and $\rho \notin \Lambda_-^{\rm ext}$ then there is $m \in \mathbb{N}$ such that $\kappa^m(\rho) = \omega$, infinite point.

After j reflections lie all reflected rays except may be the last one in some neighborhood of $(a_1, 0)$. After ϵ-small perturbation of the outgoing ray it will have the same number of reflections as an unperturbed one. Thus we have the following lemma.

Lemma 3. *For each $j = 0, 1, \dots$ there exists $\epsilon(j) > 0$ such that*

$$\operatorname{dist}(\widetilde{\Omega}_+(j+1), \mathcal{H} \setminus \widetilde{\Omega}_+(j)) > \epsilon(j) > 0.$$

Define $\Omega_+(j) := \widetilde{\Omega}_+(j) \setminus \widetilde{\Omega}_+(j+1) \subset T^*X$ such that the half-rays issued from the points of $\Omega_+(j)$ come back exactly j times and then disappears to infinity. For example $\Omega_+(1)$ corresponds to the region in $\mathcal{H}$ such that if $\rho \in \Omega_+(1)$ than $\kappa(\rho) = \rho' \in T^*X$ but $\kappa(\rho') = \omega$ – infinite point (only one reflection). We have

$$\Omega_+(0) := \mathcal{H} \cup \mathcal{G} \setminus D(\kappa),\ \Omega_+(1) := \kappa^{-1}(\Omega_+(0)),\ \dots$$
$$\mathcal{H} \cup \mathcal{G} = \bigcup_{j=0}^{\infty} \Omega_+(j),\ \kappa:\ \Omega_+(j) \mapsto \Omega_+(j-1).$$

Lemma 3 shows that approaching the glancing surface (possibly tangentially) $\Omega_+(j)$, $j = 0, 1, \dots,$ are well separated.

We define also

$$\Omega_+(j + \frac{1}{2}) := \{\rho \in \Omega_+(j);\ \operatorname{dist}(\rho, \Omega_+(j+1) < \operatorname{dist}(\rho, \Omega_+(j-1))\},$$
$$\Omega_+(j - \frac{1}{2}) := \{\rho \in \Omega_+(j);\ \operatorname{dist}(\rho, \Omega_+(j+1)) \geq \operatorname{dist}(\rho, \Omega_+(j-1))\}.$$

Thanks to Lemma 3 these domains extend naturally to $\mathcal{E}$. We denote such extended domains again by $\Omega_+(j)$.

Let G_+^0 be a step function which is constant on each $\Omega_+(j),\ j = 0, \frac{1}{2}, 1, \frac{3}{2}, \dots, N$, defined as follows

$$G^0_{+|\cup_{j\geq N}\Omega_+(j)} = 0,\ G^0_{+|\Omega_+(N-1/2)} = \frac{1}{4} h \ln^2(1/h),$$
$$G^0_{+|\Omega_+(N-1)} = \frac{1}{2} h \ln^2(1/h),\ \dots,\ G^0_{+|\Omega_+(0)} = N \cdot \frac{1}{2} h \ln^2(1/h)$$
$$G^0_{+|\mathcal{E}} = 0$$

Let $\omega \in C_0^\infty(\mathbb{R}^n)$ be such that $\omega(x) \geq 0$, $\omega(x) = 0$, for $|x| > 1$, $\int_{\mathbb{R}^n} \omega(x)dx = 1$. Put $\omega_\epsilon(x) = \epsilon^{-n}\omega\left(x/\epsilon\right)$. For $u \in C^\infty(\mathbb{R}^n)$ we consider convolution

$$(\omega_\epsilon * u)(x) = \int \omega_\epsilon(y)u(x-y)dy = \int \omega(y)u(x-\epsilon y)dy.$$

The mollifier operator $u \mapsto \omega_\epsilon * u$ extends to the continuous linear operator $D' \mapsto C^\infty$ with

$$\operatorname{supp}(\omega_\epsilon * u) \subset \operatorname{supp} u + \{x \in \mathbb{R}^n : \ |x| \leq C\epsilon\}$$

for some $C > 0$.

Using the partition of identity on T^*X and mollifiers on any coordinate neighborhood we can construct the regularization $G_+ \in C^\infty(T^*X)$ such that for $\rho \in \bigcup_{j=1}^N \Omega_+(j)$ and ϵ small enough

$$G_+(\kappa(\rho)) - G_+(\rho) \geq \frac{1}{4}h(\ln^2(1/h)) > 0.$$

In the similar way we define G_- associated to $\Omega_-(j)$ defined as before but for the incoming half-rays (by replacing κ with κ^{-1}):

$$\begin{aligned}
&\widetilde{\Omega}_-(0) = \overline{\mathcal{H}} = \mathcal{H} \cup \mathcal{G},\ \widetilde{\Omega}_-(1) = D(\kappa^{-1}) = R(\kappa),\ \text{for } j \geq 1,\ \widetilde{\Omega}_-(j) := D(\kappa^{-j}) = R(\kappa^j),\\
&\widetilde{\Omega}_-(0) \supset \widetilde{\Omega}_-(1) \supset \widetilde{\Omega}_-(2) \supset \dots,\ \bigcap_{j=0}^{\infty} \widetilde{\Omega}_-(j) = \Lambda_+^{\mathrm{ext}},\\
&\Lambda_+^{\mathrm{ext}} = \{(x,\xi) \in T^*X,\ \forall j \in \mathbb{N},\ \exists \kappa^{-j}(x,\xi) \in T^*X\},
\end{aligned}$$

where Λ_+^{ext} denotes extension of Λ_+ by the iterated action of κ^{-1}. If $\rho \notin \Lambda_+^{\mathrm{ext}}$ then there is $m \in \mathbb{N}$ such that $\kappa^{-m}(\rho) = \omega$. We have an analogue of Lemma 3.

Define $\Omega_-(j) := \widetilde{\Omega}_-(j) \setminus \widetilde{\Omega}_-(j+1)$. We have

$$\begin{aligned}
&\Omega_-(0) := \mathcal{H} \cup \mathcal{G} \setminus D(\kappa^{-1}),\ \Omega_-(1) := \kappa(\Omega_-(0)),\ \dots\\
&\mathcal{H} \cup \mathcal{G} = \bigcup_{j=0}^{\infty} \Omega_-(j),\ \kappa^{-1} :\ \Omega_-(j) \mapsto \Omega_-(j-1).
\end{aligned}$$

$$\begin{aligned}
&\Omega_-(j+\frac{1}{2}) := \{\rho \in \Omega_-(j);\ \operatorname{dist}(\rho, \Omega_-(j+1) < \operatorname{dist}(\rho, \Omega_-(j-1)\},\\
&\Omega_-(j-\frac{1}{2}) := \{\rho \in \Omega_-(j);\ \operatorname{dist}(\rho, \Omega_-(j+1)) \geq \operatorname{dist}(\rho, \Omega_-(j-1))\}.
\end{aligned}$$

We denote the extension of $\Omega_-(j)$ to $\mathcal{E}$ again by $\Omega_-(j)$.

Let G_-^0 be the step function:

$$\begin{aligned}
&G^0_{-|\cup_{j\geq N}\Omega_+-(j)} = 0,\ G^0_{-|\Omega_-(N-1/2)} = \frac{1}{4}h\ln^2(1/h),\\
&G^0_{-|\Omega_-(N-1)} = \frac{1}{2}h\ln^2(1/h),\ \dots,\ G^0_{-|\Omega_-(0)} = N \cdot \frac{1}{2}h\ln^2(1/h),\\
&G^0_{-|\mathcal{E}} = 0.
\end{aligned}$$

We define the regularization $G_- \in C^\infty(T^*X)$ such that

$$\forall\ \rho \in \bigcup_{j=1}^{N} \Omega_-(j),\ G_-(\kappa^{-1}(\rho)) - G_-(\rho) \geq \frac{1}{4} h \ln^2(1/h) > 0\ \Leftrightarrow$$

$$\Leftrightarrow G_-(\kappa(\rho')) - G_-(\rho') \leq -\frac{1}{4} h \ln^2(1/h) < 0,\ \rho' = \kappa^{-1}(\rho) \in \bigcup_{j=0}^{N-1} \Omega_-(j).$$

Let $G_{\text{ext}} := G_+ - G_-$.

We define

$$W_2 = W_2(N) := \bigcup_{i=1}^{N} \Omega_+(i) \cup \bigcup_{j=0}^{N-1} \Omega_-(j). \tag{25}$$

Lemma 4. *For all $\rho \in W_2(N)$ we have*

$$G_{\text{ext}}(\kappa(\rho)) - G_{\text{ext}}(\rho) \geq \frac{1}{2} h \ln^2(1/h).$$

Proof: If $\rho \in \Omega_+(i)$, $i \in [1, N]$, then $\exists\ j \in [0, N-1]$ such that $\rho \in \Omega_-(j)$, then we have

$$G_{\text{ext}}(\kappa(\rho)) - G_{\text{ext}}(\rho) = (G_+(\kappa(\rho)) - G_+(\rho)) - (G_-(\kappa(\rho)) - G_-(\rho)) \geq$$
$$\geq \frac{1}{4} h \ln^2(1/h) - (-\frac{1}{4} h \ln^2(1/h)).$$

∎

Thus we gain at least $\frac{1}{2} h \ln^2(1/h)$ on W_2.

We have that $\Lambda_-^{\text{ext}} \cup \bigcup_{i=N+1}^{\infty} \Omega_+(i) = \Omega_+(N+1)$ is a neighborhood of Λ_-^{ext}, and $\Lambda_+^{\text{ext}} \cup \bigcup_{j=N}^{\infty} \Omega_-(j) = \Omega_-(N)$ is a neighborhood of Λ_+^{ext}.

Let

$$W_1 = \Omega_+(N+1) \cap \Omega_-(N). \tag{26}$$

If

$$\rho \in \Lambda_-^{\text{ext}} \cup \bigcup_{i=N+1}^{\infty} \Omega_+(i) \setminus W_1 = \Omega_+(N+1) \setminus (\Omega_+(N+1) \cap \Omega_-(N)) = \text{neigh}(\Lambda_-^{\text{ext}}) \setminus W_1$$

then $G_+(\rho) = 0$ and there is j such that $\rho \in \bigcup_{j=0}^{N-1} \Omega_-(j)$ and $G_{\text{ext}}(\kappa(\rho)) - G_{\text{ext}}(\rho) = -G_-(\kappa(\rho)) + G_-(\rho) \geq \frac{1}{4} h \ln^2(1/h)$. Similar for $\rho \in \Lambda_+^{\text{ext}} \cup \bigcup_{j=N}^{\infty} \Omega_-(j) \setminus W_1$. Thus we gain there too at least $\frac{1}{4} h \ln^2(1/h)$.

Together with Lemma 4 we get that on $\overline{\mathcal{H}} \setminus W_1$ we have $G_{\text{ext}}(\kappa(\rho)) - G_{\text{ext}}(\rho) \geq \frac{1}{4} h \ln^2(1/h)$.

Inside W_1 we use $G_{\text{int}} := G_1 \circ f^{-1}$, the interior escape function defined in the previous section, and apply Lemma 2.

Let C be a large constant and $\chi \in C_0^\infty(T^*X)$ such that $\chi = 1$ in W_1. We define the global escape function

$$G(\rho) = \frac{1}{C} \chi(\rho) G_{\text{int}}(\rho) + G_{\text{ext}}(\rho) \in C^\infty(T^*X).$$

Then

$$G(\kappa(\rho)) = \frac{1}{C} \chi(\kappa(\rho)) G_{\text{int}}(\kappa(\rho)) + G_{\text{ext}}(\kappa(\rho))$$

and $G(\kappa(\rho)) - G(\rho)$ verifies the inequalities stated in Lemma 1. ∎

4. The FBI transform of Helffer and Sjöstrand and the Bargman transform with the global choice of phase.

In this section we remind the definition of the space $H(\Lambda_{tG})$ as in [10]. We follow the presentation in [23].

4.1. **Complex domains and symbol classes: generalities.** Let $\tilde{X}$ denote a complex neighborhood of X. Let $\Lambda \in T^*\widetilde{X}$ be a closed I–Lagrangian manifold which is close to T^*X in the C^∞–sense and which coincides with this set outside a compact set. Recall that "I–Lagrangian" means Lagrangian for the real symplectic form $\operatorname{Im}\sigma$, where $\sigma = \sum d\alpha_{\xi_j} \wedge d\alpha_{x_j}$ is the standard complex symplectic form. This means that if we choose (analytic) coordinates y in X and let (y,η) be the corresponding canonical coordinates on T^*X and $T^*\widetilde{X}$, then Λ is of the form $\{(y,\eta)+iH_{tG(y,\eta)};\ (y,\eta)\in T^*X\}$ for some real-valued smooth function $tG(y,\eta)$ which is close to 0 in the C^∞–sense and has compact support in η.

Here H_G denotes the Hamilton field of G. Since Λ is close to T^*X, it is also R–symplectic in the sense that the restriction to Λ of $\operatorname{Re}\sigma$ is non-degenerate. We say that Λ is an IR–manifold.

A smooth function $a(x,\xi;h)$, defined on Λ or on a suitable neighborhood of T^*X in $T^*\widetilde{X}$ is said to be of class $S^{m,k}$, if $\partial_x^p\partial_\xi^q a = \mathcal{O}(1)h^{-m}\langle\xi\rangle^{k-|q|}$.

A formal classical symbol $a \in S^{m,k}_{\mathrm{cl}}$ is of the form $a \sim h^{-m}(a_0 + ha_1 + \ldots)$ where $a_j \in S^{0,k-j}$ is independent of h. Here and in the following, we let $0 < h \leq h_0$ for some sufficiently small $h_0 > 0$. When the domain of definition is real or equal to Λ, we can find a realization of a in $S^{m,k}$ (denoted by the same letter a) so that

$$a - h^{-m}\sum_0^N h^j a_j \in S^{-(N+1)+m,k-(N+1)}.$$

When the domain of definition is a complex domain, we say that $a \in S^{m,k}_{\mathrm{cl}}$ is a formal classical analytic symbol ($a \in S^{m,k}_{\mathrm{cla}}$) if a_j are holomorphic and satisfy $|a_j| \leq C_0 C^j (j!)|\langle\xi\rangle|^{k-j}$.

It is then standard, that we can find a realization $a \in S^{m,k}$ (denoted by the same letter a) such that

$$\partial_x^k\partial_\xi^l\overline{\partial}_{x,\xi}a = \mathcal{O}(1)e^{-|\langle\xi\rangle|/Ch},\quad \left|a - h^{-m}\sum_{0\leq j\leq|\langle\xi\rangle|/C_0h} h^j a_j\right| \leq \mathcal{O}(1)e^{-|\langle\xi\rangle|/C_1h},$$

where in the last estimate $C_0 > 0$ is sufficiently large and $C, C_1 > 0$ depend on C_0. We will denote by $S^{m,k}_{\mathrm{cl}}$ and $S^{m,k}_{\mathrm{cla}}$ also the classes of realizations of classical symbols. We say that a classical (analytic) symbol $a \sim h^{-m}(a_0 + ha_1 + \ldots)$ is elliptic, if a_0 is elliptic, so that $a_0^{-1} \in S^{0,-k}$.

4.2. **The FBI transforms.** Let $X = \partial\Omega_1$. Following Sjöstrand [23] and Zworski [29] we introduce an FBI transform which is a map $T^{HS} : C^\infty(X) \mapsto C^\infty(T^*X)$ given by

$$T^{HS}u(\alpha,h) = \int_X e^{i\phi(\alpha,x)/h}a(\alpha,x;h)\chi(\alpha,x)u(x)dx, \tag{27}$$

where χ is smooth cut-off with support close to the diagonal

$$\Delta = \{(\alpha,x) \in T^*X \times X,\ \alpha_x = x\}$$

and equal to 1 in a neighborhood of Δ, $a \in S^{\frac{3n}{4},\frac{n}{4}}$ is elliptic, and ϕ is an admissible phase function. There exists $b(\alpha,x;h) \in S^{\frac{3n}{4},\frac{n}{4}}$ such that if

$$S^{HS}v(x,h) = \int_{T^*X} e^{-i\phi^*(\alpha,x)/h}b(\alpha,x;h)\chi(\alpha,x)v(x)dx,\ \ \phi^*(\alpha,x) = \overline{\phi(\overline{x},\overline{\alpha})}, \tag{28}$$

then $S^{HS}T^{HS}u = u + Ru$, where R has a distributional kernel $R(x, y; h)$ satisfying

$$|\partial_x^k \partial_y^l R| \leq C_{k,l} e^{-1/c_0 h}.$$

Let $G \in C^\infty(T^*X)$ be the escape function constructed in the previous sections. We define its C^∞ extension to $\widetilde{T^*X}$, the complex neighborhood of T^*X, with the property that

$$(dG)_\rho|_{JT_\rho(T^*X)} = 0,$$

where $J: \ T_\rho(\widetilde{T^*X}) \mapsto T_\rho(\widetilde{T^*X})$ is the complex involution. Denoting the extension by the same symbol, $G \in C^\infty(\widetilde{T^*X})$, we now define a C^∞ $I-$lagrangian, $R-$symplectic submanifold of $\widetilde{T^*X}$:

$$\Lambda_{tG} = \exp(tH_G^{\operatorname{Im}\sigma})(T^*X) \subset \widetilde{T^*X}, \ \operatorname{Im}\sigma(x, H_G^{\operatorname{Im}\sigma}) = dG(x), \ \sigma = d\alpha_\zeta \wedge d\alpha_z. \tag{29}$$

Here by a C^∞ manifold we mean a graph of a C^∞ function. We remark that a C^∞ graph in $\widetilde{T^*X}$ over T^*X which is $I-$lagrangian (i.e. its almost everywhere defined tangent plane is Lagrangian with respect to $\operatorname{Im}\sigma$) can be locally written as a graph of a differential of a C^∞ function, $\widetilde{G}$ on $T^*X \simeq T^*\mathbb{R}^n$ (locally):

$$\{\alpha = (\alpha_x, \alpha_\xi) \in T^*\mathbb{C}^n : \ \operatorname{Im}\alpha_x = \frac{\partial \widetilde{G}}{\partial \operatorname{Re}\alpha_\xi}(\operatorname{Re}\alpha_x, \operatorname{Re}\alpha_\xi), \ \operatorname{Im}\alpha_\xi = -\frac{\partial \widetilde{G}}{\partial \operatorname{Re}\alpha_x}(\operatorname{Re}\alpha_x, \operatorname{Re}\alpha_\xi)\},$$

$$(\operatorname{Re}\alpha_x, \operatorname{Re}\alpha_\xi) \in T^*\mathbb{R}^n \simeq T^*X, \ \widetilde{G} = tG.$$

The form $-\operatorname{Im}\zeta dz|_{\Lambda_{tG}}$ is (formally) closed and, as Λ_{tG} is close to T^*X, it has a primitive which is a C^∞ function on Λ_{tG}. We denote it by $H \in C^\infty(\Lambda_{tG}; \mathbb{R})$. We normalize H by demanding that it is equal to 0 for $|\alpha_\zeta|$ large enough.

Parametrizing Λ_{tG} by T^*X :

$$\Lambda_{tG} = \{(x + it\frac{\partial G}{\partial \xi}, \xi - it\frac{\partial G}{\partial x}); \ (x, \xi) \in T^*X\},$$

we get

$$H_t(\alpha_x, \alpha_\xi) = -\xi \cdot t\partial_\xi G(x, \xi) + tG(x, \xi), \ (\alpha_x, \alpha_\xi) = (x, \xi) + itH_G(x, \xi), \ (x, \xi) \in T^*X. \tag{30}$$

The weight function $H_t(\alpha_x, \alpha_\xi)$ have the same properties as tG due to Proposition 1. Thanks to the global properties of the amplitude and the phase function of T^{HS} we can continue it in α to a neighborhood of T^*X, $\widetilde{T^*X}$, in $T^*\tilde{X}$ and in particular we can define

$$T_{\Lambda_{tG}}u(\alpha; h) = Tu|_{\Lambda_{tG}}(\alpha; h), \ \alpha \in \Lambda_{tG}.$$

A deformation argument as in Sjöstrand gives an approximate inverse $S_{\Lambda_{tG}}$, with the same properties as S above, defined by (28) with T^*X replaced by Λ_{tG}. We put for $u \in C^\infty$

$$\|u\|_{tG}^2 = \|u\|^2_{H(\Lambda_{tG}, \langle\alpha_\xi\rangle^m)} = \|T_{\Lambda_{tG}}u\|^2_{L^2(\Lambda_{tG}, \langle\alpha_\xi\rangle^m e^{-H_t/h})} = \int_{\Lambda_{tG}} |T_{\Lambda_{tG}}u(\alpha; h)|^2 |\langle\alpha_\xi\rangle|^{2m} e^{-2H_t(\alpha)/h} d\alpha. \tag{31}$$

Using $S_{\Lambda_{tG}}$ we can show the independence of this norm of the choice of a specific phase function.

From [23] and [29] it follows also that the kernel of $T_{\Lambda_{tG}}S_{\Lambda_{tG}}$ has the form

$$T_{\Lambda_{tG}}S_{\Lambda_{tG}}(\alpha, \beta) = c(\alpha, \beta)e^{-\frac{i}{h}\psi(\alpha, \beta)} + r(\alpha, \beta; h),$$

where $\psi(\alpha, \beta) = v.c._z(\phi(\alpha, z) - \phi^*(\beta, z))$, $c(\alpha, \beta; h) = \mathcal{O}(h^{-n})$ is supported near $\alpha = \beta$,

$$r(\alpha, \beta; h) = \mathcal{O}_N(1)e^{-1/Ch}|\langle\beta_\zeta\rangle|^{-N}$$

and

$$-H(\alpha) - \operatorname{Im}\psi(\alpha, \beta) + H(\beta) \leq -\frac{1}{c}d(\alpha, \beta)^2.$$

Here $d(\alpha,\beta)$ is any non-degenerate distance on Λ_{tG}. We used that $|\nabla^2 G| = \mathcal{O}(1)$.

4.3. Generalities on the Bargman transforms with the global choice of phase and choice of the norm.

Following [23] we replace the norm $\|\cdot\|_{tG}$ by an equivalent norm obtained by decomposition of T^*X.

In the region where ξ is bounded, it will be convenient to work with transforms which are holomorphic up to exponentially small errors, and for that we make a different choice of T, and take an FBI–transform of Bargman type with a global choice of phase.

The standard Bargman transform in $\mathbb{R}^n$ is given by

$$T_0u(z,h) = h^{-\frac{3n}{4}}\int e^{-(z-x)^2/2h}u(x)dx,\ T_0 : L^2(\mathbb{R}^n) \mapsto H_{\Phi_0}(\mathbb{C}^n),\ \Phi_0(z) = (\operatorname{Im} z)^2/2,$$

where $H_\Phi := \{u \text{ holomorphic},\ \int |u|^2 e^{-2\Phi/h}dx < \infty\}$, associated to the complex canonical transformation

$$\kappa_{T_0}(x,\xi) = (x - i\xi, \xi),\ \kappa(\mathbb{R}^{2n}) = \Lambda_{\Phi_0} := \{z = \frac{2}{i}\frac{\partial \Phi_0(z)}{\partial z}\}.$$

Let $\phi_0(\alpha,x) = i(\operatorname{Im} z)^2/2 + i(z-x)^2/2$, $z = \alpha_x - i\alpha_\xi$. Then $\phi_0(\alpha,x)$ is admissible phase function and thus $e^{-2\Phi_0(z)/h}T_0$ is local FBI transform of the type described in the previous section:

$$e^{-2\Phi_0(z)/h}T_0u(\alpha) = h^{-\frac{3n}{4}}\int e^{\frac{i}{h}((\alpha_x - x)\alpha_\xi + i(x-\alpha_x)^2/2)}u(x)dx.$$

$$\kappa_{T_0}(\Lambda_{tG}) = \Lambda_{\Phi_t},\ \Phi_t(z) = \Phi_0(z) + tG(\operatorname{Re} z, -\operatorname{Im} z) + \mathcal{O}(t^2).$$

We need the global version of Bargman transform on T^*X. We equip X with some analytic Riemannian metric so that we have a distance dist and a volume density dy.

The function $\operatorname{dist}(x,y)^2$ is analytic in a neighborhood of the diagonal in $X \times X$, so we can consider it as a holomorphic function in a region

$$\left\{(x,y) \in \widetilde{X} \times \widetilde{X};\ \operatorname{dist}(x,y) < \frac{1}{C},\ |\operatorname{Im} x|, |\operatorname{Im} y| < \frac{1}{C}\right\}.$$

Put

$$\phi(x,y) = i\lambda_0 \operatorname{dist}(x,y)^2, \tag{32}$$

where $\lambda_0 > 0$ is a constant that we choose large enough, depending on the size of the neighborhood of the zero section in T^*X, that we wish to cover.

For $x \in \widetilde{X}$, $|\operatorname{Im} x| < 1/C$, put

$$Tu(x;h) = h^{\frac{-3n}{4}}\int e^{\frac{i}{h}\phi(x,y)}\chi(x,y)u(y)dy,\ u \in \mathcal{D}'(X), \tag{33}$$

where χ is a smooth cut-off function with support in

$$\{(x,y) \in \widetilde{X} \times X;\ |\operatorname{Im} x| < 1/C,\ \operatorname{dist}(y, y(x)) < 1/C\}$$

containing $\mathcal{H} \cup \mathcal{G}$. Here $y(x) \in X$ is the point close to x, where $X \ni y \mapsto -\operatorname{Im} \phi(x,y)$ attains its non-degenerate maximum. We collect the following facts from [23]:

- The function $\Phi_0(x) = -\operatorname{Im} \phi(x, y(x))$, $x \in \widetilde{X}$, $|\operatorname{Im} x| < 1/C$, is strictly plurisubharmonic and is of the order of magnitude $\sim |\operatorname{Im} x|^2$.
- $\Lambda_{\Phi_0} := \{(x, \frac{2}{i}\partial\Phi_0) \in T^*\widetilde{X}\}$ is an IR–manifold given by $\Lambda_{\Phi_0} = \kappa_T(T^*X)$, where κ_T is the complex canonical transform associated to T, given by $(y, -\phi'_y(x,y)) \mapsto (x, \phi'_x(x,y))$. Here and in the following, we identify $\widetilde{X}$ with its intersection with a tubular neighborhood of X which is independent of the choice of λ_0 in (32).

- If $L^2_{\Phi_0} = L^2(\widetilde{X};\ e^{-2\Phi_0/h}L(dx))$, for $L(dx)$ denoting a choice of Lebesgue measure (up to a non-vanishing continuous factor), then $T = \mathcal{O}(1):\ L^2(X) \mapsto L^2_{\Phi_0}$, $\overline{\partial}_x T = \mathcal{O}(e^{-1/Ch}):\ L^2(X) \mapsto L^2_{\Phi_0}$. This means that up to an exponentially small error Tu is holomorphic for $u \in L^2(X)$ (and even for $u \in \mathcal{D}'(X)$).
- Unitarity: Modulo exponentially small errors and microlocally, T is unitary
$$L^2(X) \mapsto L^2(\widetilde{X}; a_0 e^{-2\Phi_0/h}L(dx)),$$
where $L(dx)$ is chosen as indicated above, and $a_0(x;h)$ is a positive elliptic analytic symbol of order 0.
- Let $\Lambda_{tG} \subset T^*\widetilde{X}$ be an IR–manifold as before. Then $\kappa_T(\Lambda_{tG}) = \Lambda_{\Phi_t}$, where Φ_t can be normalized by the requirement that $\Phi_t = \Phi_0$ near the boundary of $\widetilde{X}$. (Here is where we have to choose λ_0 large enough, depending on Λ_{tG}. In the applications, for a given elliptic operator, Λ_{tG} and T^*X will coincide outside a fixed compact neighborhood of the zero section, and the whole study will be carried out with a fixed λ_0.)

Locally we can identify $T^*X \simeq T^*\mathbb{R}^n$ and take $\phi(x,y) = i(x-y)^2/2$. Then locally $\kappa_T :$ $(y,\eta) \mapsto (y - i\eta, \eta) =: \Lambda_{\Phi_0}$ with $\Phi_0 = (\operatorname{Im} x)^2/2$. Using the local formula
$$H_t(y,\eta) = -\operatorname{Re}\eta \cdot \operatorname{Im} y + tG(\operatorname{Re} y, \operatorname{Re}\eta)$$
we get
$$\begin{aligned}
&\Phi_t(x) = v.c._{(y,\operatorname{Re}\eta)\in\mathbb{C}^n\times\mathbb{R}^n}\,(-\operatorname{Im}\phi(x,y) - \operatorname{Re}\eta\cdot\operatorname{Im} y + tG(\operatorname{Re} y,\operatorname{Re}\eta)) \ \Leftrightarrow\\
&\Phi_t(x) = \Phi_0(x) + tG(x) + \mathcal{O}(t^2|x|^2),\\
&\text{where } G(x) = G(y(x),\eta(x)),\ (y(x),\eta(x)) = \kappa_T^{-1}\left(x, \frac{2}{i}\frac{\partial\Phi_0(x)}{\partial x}\right).
\end{aligned}\tag{34}$$

Let W be a neighborhood of $\rho_1 \in T^*X$, such that $W_1 \subset\subset W$. We identify W with $\kappa_T(W)$.

It follows from [10], Section 9, that for $u \in H(\Lambda; \langle\alpha_\xi\rangle^m)$ we have
$$\|Tu\|_{H_{\Phi_t}(W)} \le c\|u\|_{tG}.$$
Let $W' \subset\subset W$ and $W_1 \subset\subset W'$ be another neighborhood of a_1 which is identified with $\{\alpha \in \Lambda_{tG};\ \operatorname{Re}\alpha \in W'\}$. One can show, using methods from [10], [22], that for $0 \le t < t_0$ with t_0 small enough $\exists\ \delta_0 > 0$ such that
$$\|u\|_{tG,W'} \le \mathcal{O}(1)\left(\|Tu\|_{H_{\Phi_t}(W)} + e^{-\delta_0/h}\|u\|_{tG,\complement W'}\right),$$
where in general the seminorm $\|u\|_{tG,W}$ is defined as in (31) by integration over $W \subset \Lambda_{tG}$ only.

We define a new norm uniformly equivalent to $\|\cdot\|_{tG}$ as $h \to 0$ by
$$\|u\|_{tG,\complement W'} + \|Tu\|_{H_{\Phi_t}(W)}.$$

5. The FBI-Bargman transform of the quantum billiard operator M.

The operator M was defined in the Introduction, formula (12). Here we give an alternative definition (see Burq [4]).

Let $h = 1/\lambda \in \mathbb{R}_+$. If $H_{1,+}$ is the outgoing resolvent of the problem
$$\begin{cases} (-h^2\Delta - 1)H_{1,+}u &= 0, \text{ in } \complement\Omega_1\\ H_{1,+}u_{|\partial\Omega_1} &= u, \end{cases}\tag{35}$$
we consider the outgoing resolvent of the problem
$$\begin{cases} (-h^2\Delta - 1)H_{21}u &= 0, \text{ in } \complement\Omega_2\\ H_{21}u_{|\partial\Omega_2} &= H_{1,+}u_{|\partial\Omega_2}. \end{cases}\tag{36}$$

The operator M is defined as

$$Mu = H_{21} u_{|\partial\Omega_1}. \tag{37}$$

Some known facts about M are collected in Appendix A.

Let W_1 be the neighborhood of $\rho_1 = (a_1, 0) \in T^*X$ independent of h and defined in (26): $W_1 = \Omega_+(N+1) \cap \Omega_-(N)$. Let G be the global escape function as in Proposition 1.

5.1. **Properties of the kernel of M on $\complement W_1$.**

Let T be the FBI transform of Helffer and Sjöstrand as in (27) with admissible phase function ϕ in a complex neighborhood $U \times V$ of (ρ_1, a_1) and let S be an approximate inverse to T as in (28). Let $\mu, \rho \in T^*X$ be arbitrary real for the moment. As M is bounded operator $L^2(X) \mapsto L^2(X)$ and T, S are bounded $L^2(X) \mapsto L^2(T^*X)$ respectively $L^2(T^*X) \mapsto L^2(X)$ then the distributional kernel K of TMS given by

$$TMSTu(\alpha, h) = \int_{\mathrm{neigh}\,(\rho_1)} K(\alpha, \beta; h) Tu(\beta) d\beta, \ \alpha \in \mathrm{neigh}\,(\rho_1), \tag{38}$$

is well defined.

By the results on the propagation of Gevrey 3 singularities due to Lebeau [16] (see also Burq [4]) it is known that on the complement of the graph (κ) :

$$\{(\alpha, \beta) \in T^*X \times T^*X, \ \alpha = \kappa(\beta)\}, \tag{39}$$

$K(\alpha, \beta; h)$ is small:

$$K(\alpha, \beta; h) = \mathcal{O}_N(1) e^{-1/ch^{1/3}} (\max(|\langle \alpha_\xi \rangle|, |\langle \beta_\xi \rangle|))^{-N} \ \text{ if } \alpha \neq \kappa(\beta), \ \text{for any } N > 0, \tag{40}$$

if $\kappa(\rho)$ is defined, otherwise no condition. Since $H(\alpha)$ has compact support and is small, we have the same estimate for the reduced kernel: $e^{(-H(\alpha)+H(\beta))/h} K(\alpha, \beta; h)$.

In a neighborhood $U \times U$ of the graph of κ (39), from the general principles, as M is bounded, we know only that $K(\alpha, \beta; h)$ is of order $\mathcal{O}(h^{-N'})$ for some $N' > 0$. If $\alpha, \beta \in \complement W_1$ we can get a better bound on the reduced kernel $e^{(-H(\alpha)+H(\beta))/h} K(\alpha, \beta; h)$ by using the properties of the escape function G, Lemma 1: if $\alpha = \kappa(\beta) \in T^*X \setminus W_1$ then $G(\alpha) - G(\beta) \geq Ch \ln^2(1/h)$ and the bound is still valid in the whole neighborhood of $\{(\alpha, \beta) \in T^*X \setminus W_1; \ \alpha = \kappa(\beta)\}$.

Let Λ_{tG} be the IR-submanifold of the complex neighborhood of T^*X defined in (29). Let $K(\alpha, \beta; h)$ denote also the extension of (38) to the complex neighborhood of (ρ_1, ρ_1) of the form $(U \cap \Lambda_{tG}) \times (U \cap \Lambda_{tG})$. Let $H \in C^\infty(\Lambda_{tG}; \mathbb{R})$ satisfy $d_\alpha H_{|\Lambda_{tG}} = -\mathrm{Im}\, \alpha_\xi d\alpha_{x|\Lambda_{tG}}$ and be equal to zero for $|\alpha_\xi|$ large enough. By (30) we have locally

$$H_t(\alpha_x, \alpha_\xi) = -\mathrm{Re}\, \alpha_\xi \cdot \mathrm{Im}\, \alpha_\xi + tG(\mathrm{Re}\, \alpha_x, \mathrm{Re}\, \alpha_\xi).$$

Then for (α, β) in a neighborhood of $\{(\alpha, \beta) \in \complement W_1; \ \mathrm{Re}\,(\alpha) = \kappa(\mathrm{Re}\, \beta)\}$ we can arrange that

$$e^{-H(\alpha)/h} K(\alpha, \beta; h) e^{H(\beta)/h} = e^{-\frac{1}{Ch} \min(h \ln^2(1/h), |\alpha - \kappa(\beta)|^2)} (\max(|\langle \alpha_\xi \rangle|, |\langle \beta_\xi \rangle|))^{-N} \tag{41}$$

for some $N_0 > 0$ and for any $N > 0$.

Then by the general theory of [10] and [23] the estimate (41) implies that for $u \in H(\Lambda_{tG})$ we have the bound:

$$\|Mu\|_{tG, \complement W'} \leq \mathcal{O}(h^{N_0}) \|u\|_{tG} \tag{42}$$

for some $N_0 > 0$. Here W' is any $h-$independent neighborhood of ρ_1, $W_1 \subset\subset W' \subset\subset W$ identified with $\{\alpha \in \Lambda_{tG}; \ \mathrm{Re}\, \alpha \in W'\}$ and in general the seminorm $\|u\|_{tG,W}$ is defined as in (31) by integration over $W \subset \Lambda_{tG}$ only (see the end of Section 4.3). Bound (42) extends to complex $\lambda \in \Lambda_{A,B}$ with some different power $N_0 > 0$.

5.2. **Local form of M.** Microlocally near $(a_1, 0) \in T^*\partial\Omega_1$, the asymptotic solution to (36) is given by means of WKB construction.

In the hyperbolic zone $\mathcal{H}$, M is approximated by Fourier integral operator $H \in I_h^0(X, X; \kappa')$, where $\kappa' = \{(x, \xi; y, -\eta) : \ (x, \xi) = \kappa(y, \eta)\}$, associated to the real canonical transformation of billiard κ, with the real phase and can be taken in the form (see [9]):

$$Hu(u, \lambda) = \left(\frac{\lambda}{2\pi}\right)^n \int e^{-i\lambda(s(x,\theta)-y\theta+2d)} b(x, y, \theta; \lambda) u(y) dy d\theta, \tag{43}$$

where s solves the eikonal equation, $s(x, \theta) - y\theta$ parameterizes the canonical transformation of billiard κ, $s(0, 0) = 0$.

If λ is complex, $\lambda \in \Lambda_{A,B}$, then we denote $\lambda_1 = \operatorname{Re}\lambda$, $\lambda_2 = \operatorname{Im}\lambda$ and write

$$Hu(x, \lambda) = \left(\frac{\lambda_1}{2\pi}\right)^n \iint e^{-i\lambda_1(s(x,\theta)-y\theta+2d)} \tilde{b}(x, y, \theta; \lambda_1) u(y, \lambda) dy d\theta$$

with

$$\tilde{b}(x, y, \theta; \lambda) = (1 + i \tan \arg \lambda)^n e^{\lambda_2(s(x,\theta)-y\theta+2d)} b(x, y, \theta; \lambda_1).$$

Let $T : L^2(X) \mapsto H_{\Phi_0}$, $H(\Lambda_t) \mapsto H_{\Phi_t}$ (modulo exponentially small errors) be as in (33) associated to the complex canonical transformation κ_T such that $\kappa_T(T^*X) = \Lambda_{\Phi_0}$ and $\kappa_T(\Lambda_{tG}) \mapsto \Lambda_{\Phi_t}$.

For Φ denoting either Φ_0 or Φ_t, the Fourier integral operator $\hat{H} : H_\Phi \mapsto H_\Phi$ is associated to the real canonical transformation $\hat{\kappa}$ in the sense that locally $\hat{\kappa}(\Lambda_\Phi) = \Lambda_\Phi$.

Let $V_1 = \pi_x \kappa_T W_1$ and for any $x \in V_1$ let $\gamma(x) \in \Lambda_{tG}$ be such that $\pi_x \kappa_T(\gamma(x)) = x$. In $W_1 \subset \mathcal{H}$ we apply the Bargman transform and denote $\hat{H}$ the transformed operator associated to the canonical transformation $\hat{\kappa} = \kappa_T \circ \kappa \circ \kappa_T^{-1}$, satisfying for $u \in L^2(W_1)$,

$$TMu = \hat{H}Tu + \mathcal{O}(h^\infty)\|u\| \text{ in } H_\Phi.$$

As Hu is asymptotic solution to (36) and obstacles are analytic then $\hat{H}$ can be taken analytic of the form

$$\hat{H}u(x, \lambda) = \left(\frac{\lambda}{2\pi}\right)^n \iint e^{-i\lambda\phi(x,y,\theta)} a(x, y, \theta; \lambda) u(y, \lambda) d\theta dy, \ u \in H_\Phi. \tag{44}$$

with analytic phase and an amplitude which is an analytic symbol of order 0 realized with some suitable contour $\Gamma(x) \subset W_1$ passing through the critical point, and introducing some cut-off functions.

It is well-known that, by choosing an appropriate integration contour, $\hat{H}$ is associated with a kernel $K_{\hat{H}} \in H_{\Phi(x)+\Phi(\overline{y})}$ such that, formally:

$$\hat{H}u(x, h) = \int K_{\hat{H}}(x, \overline{y}, h) u(y, h) e^{-2\Phi(y)/h} L(dy), \ u \in H_\Phi.$$

The kernel $K_{\hat{H}}$ is uniquely defined as an element of $H_{\Phi(x)+\Phi(\overline{y})}$ through the data of $\hat{H}$, but not as a function.

Denote

$$k(x, y; h) = K_{\hat{H}}(x, \overline{y}, h) e^{-2\Phi(y)/h}.$$

If $\operatorname{Re}\gamma(x) \neq \kappa(\operatorname{Re}\gamma(y)) \notin W_0$ then by the results on the propagation of analytic singularities of Lebeau [16] (see also Burq [4]) we have

$$e^{-\Phi(x)/h} k(x, y; h) e^{\Phi(y)/h} = \mathcal{O}\left(e^{-\epsilon/h}\right), \tag{45}$$

where Φ is either Φ_0 or $\Phi_t = \Phi_0(x)+tG(x)+\mathcal{O}(t^2|x|^2)$, where $G(x) := G(\operatorname{Re}\gamma(x))$, $\pi_x\kappa_T(\gamma(x)) = x$. Here we used that $G = \mathcal{O}(h\ln(1/h))$.

If $\operatorname{Re}\gamma(x) = \kappa(\operatorname{Re}\gamma(y))$ then we use the properties of the escape function:

$$-G(\kappa(\operatorname{Re}\gamma(y))) + G(\operatorname{Re}\gamma(y)) \le -Ch\ln(1/h),\ c>0,\ \text{if } \operatorname{Re}\gamma \in W_1 \setminus W_0.$$

Choosing W_0 large enough (c_0 in the definition (22) is sufficiently large) we can arrange that

$$e^{-\Phi_t(x)/h}k(x,y;h)e^{\Phi_t(y)/h} = \mathcal{O}\left(h^{N_0}\right) \text{ if } \operatorname{Re}\gamma(x) = \kappa(\operatorname{Re}\gamma(y)) \cap V_0 = \emptyset, \tag{46}$$

for some $N_0 > 0$ and $V_0 = \pi_x\kappa_T W_0$.

5.3. **The Bargman transform reducing the stable manifolds to $x = 0,\ \xi = 0$.** The outgoing Λ_+, incoming Λ_- stable manifolds for κ are lagrangian, intersecting transversally at $(0,0)$. We can introduce real symplectic coordinates (x,ξ) such that $\Lambda_+ :\ \xi = 0,\ \Lambda_- :\ x = 0$.

We consider the image of $\Lambda_\pm$ under the complex canonical transformation $\kappa_T :\ T^*X \mapsto \Lambda_{\Phi_0}$ associated to the Bargman transform T. Transformation κ_T preserves the properties of $\Lambda_\pm$. We will often write $\Lambda_\pm$ instead of $\kappa_T(\Lambda_\pm)$.

We know that the fiber $\{x = 0\}$ is a Lagrangian manifold, which is strictly negative with respect to Λ_{Φ_0}. It follows that $\{x = 0\}$ and $\kappa(\Lambda_-)$ are transversal. Then $\Lambda_\pm$ are given by $\xi = \partial_x\phi_\pm$, where $\phi_\pm$ are holomorphic, and $-\operatorname{Im}\phi_+ - \Phi_0(x) \le 0$. The image of Λ_{tG} is of the form Λ_{Φ_t}, where $\Phi_t(x) = \Phi_0(x) + tG(x) + \mathcal{O}(t^2|x|^2)$, where G is considered also as a function on $\Lambda_{\Phi_0} \simeq \mathbb{C}^n_x$ (see (34)):

$$G(x) := G \circ (\pi_x \circ \kappa_{T|T^*X})^{-1} = G(y,\eta) \text{ with } (y,\eta) \in \Lambda_{tG} \text{ given by } \pi_x(\kappa_T(y,\eta)) = x. \tag{47}$$

(If $\phi = i(x-y)^2/2$, and $(y,\eta) \in \mathbb{R}^{2n}$ (instead of complex Λ_{tG}), then $x = y - i\eta$ and $G(x) = G(\operatorname{Re}x, -\operatorname{Im}x)$.)

The strict positivity of Λ_+ with respect to Λ_{tG} then implies that

$$\operatorname{Im}\phi_+ + \Phi_t \sim |x|^2.$$

Using the strict negativity of Λ_- with respect to Λ_{tG} we have (see [25])

Lemma 5. *There is totally real linear space, $L \subset \mathbb{C}^n$, of real dimension n, such that,*

$$\Phi_t + \operatorname{Im}\phi_- \sim -|x|^2 \text{ on } L.$$

Considering still the situation after application of κ_T, we now introduce a new canonical transformation κ_F which maps Λ_+ to $\{\xi = 0\}$ and Λ_- to $\{x = 0\}$. Then it is easy to see that κ_F is given by $\kappa_F :\ (y, -\phi'_y(x,y)) \mapsto (x, \phi'_x(x,y))$, where the generating function $\phi(x,y)$ verifies: $\det\phi''_{x,y} \neq 0,\ \phi(0,y) = -\phi_-(y)$.

Let $f(x,y;h)$ be classical analytic symbol defined near $(x,y) = (0,0)$. Using the Lemma 5, we see that if $u \in H_{\Phi_t}$, then we can define $Fu \in H_{\hat\Phi_t}$, by choosing a nice contour for the integral expression,

$$Fu(x;h) = \int e^{i\phi(x,y)/h}f(x,y;h)u(y)dy.$$

Here, we use terminology of [22], $\hat\Phi_t$ is a new strictly pl.s.h. function, determined up to a constant by the relation $\Lambda_{\hat\Phi_t} = \kappa_F(\Lambda_{\Phi_t})$.

As Λ_+ is strictly positive with respect to Λ_{Φ_t} we get that $\kappa_F(\Lambda_+)$ is strictly positive with respect to $\Lambda_{\hat\Phi_t}$ and the strict positivity of $\kappa_F(\Lambda_+) :\ \xi = 0$ (that we shall from now on denote simply by "Λ_+") means that,

$$\hat\Phi_t \sim |x|^2.$$

Up to exponentially small errors (modulo equivalence in the spaces H_{Φ_t}, $H_{\hat{\Phi}_t}$,) we can invert F by an operator,

$$Gv(y;h) = \int e^{-i\phi(x,y)/h} g(x,y;h)v(x)dx.$$

The canonical transformation of billiard κ is transformed to

$$\kappa \equiv \exp H_p,\ \mathrm{D}\kappa(0,0) = \begin{pmatrix} A & 0 \\ 0 & (A^T)^{-1} \end{pmatrix},\ p = B(x,\xi)x\xi,\ B(0,0) = \mathrm{diag}(\mu_1, \dots, \mu_n),$$

where we have used the Lewis-Sternberg theorem (see [13]) and $\equiv$ denote the equivalence relation for formal Taylor series at $(0,0)$.

We denote $\widehat{W_i} := \kappa_F(W_i)$ and $\hat{V}_i := \pi(\widehat{W_i})$ for $i = 0,1$.

Composing the Bargman transform T with the integral transform F we get an operator of norm $\mathcal{O}(1)$,

$$FT:\ H(\Lambda_{tG}) \mapsto H_\Phi(V_1). \tag{48}$$

Here $t > 0$ is small and fixed, G is the escape function introduced in Section 3, Φ denotes a function having all the properties of $\hat{\Phi}_t$ and V_1 is small open neighborhood of 0 in $\mathbb{C}^n$.

The direct definition of T and F only gives that $\overline{\partial}(FT)$ is exponentially small, but we can correct this by solving a $\overline{\partial}$-problem, using the fact that Φ is strictly plurisubharmonic.

Composed operator FT has microlocal inverse SG of norm $\mathcal{O}(1)$: $H_\Phi(V_1) \mapsto H(\Lambda_{tG})$ with the properties:

- $SGFT$ is a pseudodifferential operator of order 0 adopted to Λ_{tG} in the sense of [10], which has compactly supported symbol and which realizes the identity microlocally near $\rho_1 = (a_1, 0)$.
- We have $\|FTSGu - u\|_{H_\Phi(\tilde{V}_1)} = \mathcal{O}(h^\infty)\|u\|_{H_\Phi(V_1)}$, where $\tilde{V}_1 \subset\subset V_1$ has the same properties as V_1.

We have

Theorem 3. *The transformed operator* $FTHSG = e^{-i\lambda 2d}M_1$ *is an analytic Fourier integral operator given formally by*

$$M_1u(x,\lambda) = \left(\frac{\lambda}{2\pi}\right)^n \int e^{-i\lambda(\varphi(x,\theta) - y\cdot\theta)} b(x,y,\theta;\lambda)u(y,\lambda)dyd\theta. \tag{49}$$

Here $b \in S^{0,0}_{\mathrm{cla}}$, $b(0,0;\lambda) = (\nu_1 \cdot \ldots \cdot \nu_n)^{-1/2} + O(|\lambda|^{-1})$, *where* $\nu_1, \dots, \nu_n$ *are the eigenvalues* > 1 *of* $\mathrm{D}\kappa(0,0)$. *We have* $\varphi(x,\theta) = A^{-1}x\cdot\theta + \mathcal{O}(|(x,\theta)|^3)$, *where* A *is diagonal with eigenvalues* $\nu_1, \dots, \nu_n$, $\nu_i = e^{\mu_i}$.

M_1 *can be realized as bounded operator* $H_\Phi(\Omega_1) \mapsto H_\Phi(\Omega_2)$ *with domains* $\Omega_2 \subset\subset \Omega_1 \subset\subset V_1$, *where we identify* Ω_1 *with* $\pi\kappa\left(\Lambda_\Phi \cap \pi^{-1}\Omega_1\right)$.

For any $\chi \in C_0^\infty(\mathbb{C}^n)$ *with* $\operatorname{supp}\chi \subset V_1 \setminus V_0$ *and some* $N_0 > 0$, *we have*

$$\|\chi M_1 u\|_{L^2_\Phi(\Omega_2)} = \mathcal{O}(h^{N_0})\|u\|_{L^2_\Phi(\Omega_1)}. \tag{50}$$

The form of M_1 in (49) follows as in [9]. The estimate (50) follows from (45) and (46).

Let Φ_t denote the new $\hat{\Phi}_t$ after the transformation F.

6. Semiclassical quantum Birkhoff normal form.

In this section we perform an analytic Birkhoff transform κ_r up to some fix order r,

$$\kappa_r:\ \Lambda_{\Phi_t} \mapsto \Lambda_{\hat{\Phi}_t}.$$

Let $\lambda \in \Lambda_{A,B}$ and we put $h = 1/\mathrm{Re}\,\lambda$. We need the following notion of equivalence used in papers [13], [14]. Here ρ denotes either pair (x,ξ) or triple (x,y,θ).

Definition 2. *Let $U \in I^0(X,X,\kappa')$ and $\tilde{U} \in I^0(X,X,\tilde{\kappa}')$ be two Fourier integral operators. Then $U \equiv \tilde{U}$ to the order r if $\kappa(0,0) = \tilde{\kappa}(0,0) = (0,0)$, κ, $\tilde{\kappa}$ agree to the order ρ^{2r+1} at $(0,0)$, the terms with number j, with $0 \le j \le r$, in the asymptotic expansions (in powers of h, corresponding to h^j) of $\phi, \tilde{\phi}, a, \tilde{a}$ (phase, respectively, amplitudes) agree to the order $\rho^{2(r-j)+1}$ at $(0,0,0)$.*

We write $a \equiv \mathcal{O}(|\rho|^{2r+1} + h^{r+1})$ to denote that a is equivalent to zero to the order r, and use $\equiv$ to denote the equivalence to infinite order.

The theorem proven in [13] for real hyperbolic κ says the following:

Theorem 4. *Let M_1 be a Fourier integral operator as in Theorem 3 which quantizes an analytic canonical transformation $\kappa :\ \mathrm{neigh}\,(0,\mathbb{C}^{2n}) \to \mathrm{neigh}\,(0,\mathbb{C}^{2n})$. Assume that the eigenvalues of $d\kappa(0)$ satisfy*

$$0 < \nu_n^{-1} \le \ldots \le \nu_1^{-1} < 1 < \nu_1 \le \ldots \le \nu_n.$$

Let $\mu_j = \log \nu_j$.

Then there exists a pseudodifferential operator $P(x, \lambda^{-1}D_x; h)$ with symbol $P(\rho;h) \sim p(\rho) + hp_1(\rho) + \ldots$, such that

$$M_1 \equiv e^{-i\lambda P}. \tag{51}$$

P is uniquely determined modulo "$\equiv$" and up to an integer multiple of $\frac{2\pi}{\lambda}$ by (51) and the choice of p_0 such that $p(\rho) - p_0(\rho) = \mathcal{O}(|\rho|^3)$.

Suppose

$$\sum_1^n k_j\mu_j = 0,\ k_j \in \mathbb{Z} \implies k_1 = \ldots = k_n = 0. \tag{52}$$

Then there exists elliptic Fourier integral operator B and a classical symbol of order 0,

$$F(\imath;1/\lambda) = F_0(\imath) + \lambda^{-1}F_1(\imath) + \lambda^{-2}F_2(\imath) + \ldots,\ \imath = (\imath_1,\ldots,\imath_n),\ \imath_j = \xi_j x_j,$$

$$F_0(\imath) = \sum_{j=1}^n \mu_j\imath_j + R(\imath),\ R(\imath) = \mathcal{O}(\imath^2),$$

such that

$$P \equiv B^{-1}F(I;1/\lambda)B,\ M_1 \equiv B^{-1}e^{-i\lambda F(I;1/\lambda)}B,\ I = (I_1,\ldots,I_n),\ I_j = \frac{1}{2i\lambda}(x_j\partial_{x_j} + \partial_{x_j}x_j).$$

When M_1 is microlocally unitary near $(0,0)$, $F(I;1/\lambda)$ can be chosen to be microlocally self-adjoint and unitary, respectively, at $(0,0,0)$.

Fix some order $r \ge 1$ and let

$$F^r(\imath;h) := F_0^r(\imath) + \lambda^{-1}F_1^{r-1}(\imath) + \lambda^{-2}F_2^{r-2}(\imath) + \ldots + \lambda^{-r}F_r^0,$$

where F_j^{r-j} is polynomial of degree $(r-j)$ in $\imath$ (F_r^0 is constant) such that

$$F_j(\imath) - F_j^{r-j}(\imath) = \mathcal{O}(\imath^{r-j+1}),$$

($F_j^{r-j}(\xi\cdot x)$ is resonant up to order $2(r-j)$). For any $r' > r$ the terms in expansion of $F^{r'}(\imath;1/\lambda)$ and $F^{r'}$ with the same indexes coincide, thus the coefficients in expansion $F^r(\imath;1/\lambda)$ are independent of r.

Then $F^r(I;1/\lambda)$ is *analytic* pseudodifferential operator. From the construction of Birkhoff transform B for the *Hamiltonian* it is easy to see that there exists an *analytic* Fourier integral operator B_r associated to an analytic κ_r such that

$$P - B_r^{-1}F^r(I;1/\lambda)B_r = Q^r,$$

where Q^r is pseudodifferential operator with the symbol $q^r(\rho;1/\lambda) \sim q_0(\rho)+\lambda^{-1}q_1(\rho)+\dots$, with $q_j(\rho) = \mathcal{O}(|\rho|^{2(r-j)+1} + h^{r-j+1})$, if $0 \le j \le r$ and $q_j = p_j$ for $j \ge r+1$. Here B_r^{-1} is analytic Fourier integral operator satisfying $B_r^{-1}B_r - I = \tilde{Q}^r$ and $\tilde{Q}^r$ of the same type as Q_r.

Operator $e^{-i\lambda F^r(I;1/\lambda)}$ is analytic Fourier integral operator which satisfies

$$M_1 - B_r^{-1}e^{-i\lambda F^r(I;1/\lambda)}B_r \equiv \mathcal{O}(|\rho|^{2r+1} + h^{r+1}).$$

We denote

$$M_0 := B_r M_1 B_r^{-1},\ R^r := M_0 - e^{-i\lambda F^r(I;1/\lambda)}.$$

Operators M_0 and R^r have the form (49). Let

$$R^r u(y,h) = \lambda^n \iint e^{-i\lambda\varphi^r(x,y,\theta)} b^r(x,y,\theta,\lambda)dyd\theta$$

and $b \sim \sum \lambda^j b_j$, $b_j = \mathcal{O}((|x|+|y|+|\theta|)^{2(r-j)_+ +1})$. The integration contour is chosen such that

$$-\Phi(x) + \operatorname{Im}\varphi^r(x,y,\theta) + \Phi(y) \sim -|x|^2 - |y|^2 - |\theta|^2.$$

We have

$$e^{-\Phi/h}R^r e^{\Phi/h} \equiv \mathcal{O}(|\rho|^{2r+1} + h^{r+1}). \tag{53}$$

Relation extends to $\lambda \in \Lambda_{A,B}$ with the bound $\mathcal{O}\left(h^{-C}(|\rho|^{2r+1} + h^{r+1})\right)$ for some $C > 0$ and usual convention $h = 1/\lambda_1$, $\lambda_1 = \operatorname{Re}\lambda$.

Using (53) and that for $\lambda \in \Lambda_{A,B}$, we have $|e^{-i2d\lambda}| \le \mathcal{O}(1)\lambda_1^m$ for some (positive) m, we get

Lemma 6. *Let $W_0 = B(0, c_0\sqrt{s})$, $s = h\ln(1/h)$, $c_0 > 0$ as in (22) and V_0 such that $\pi(W_0) = V_0$. Suppose $\lambda \in \Lambda_{A,B}$. For any $N \in \mathbb{N}$ there is $r = r(N) \in \mathbb{N}$ such that*

$$\|e^{-i2d\lambda}R^r u\|_{L^2_\Phi(\Omega_2)} \le \mathcal{O}_r(h^N)\|u\|_{L^2_\Phi(\Omega_1)},$$

where $\Omega_2 \subset\subset \Omega_1 \subset\subset V_0$ and Φ is either Φ_0 or Φ_t or any pl.s.h. function close to Φ_0 in C^2.

Proof: In the proof $h = 1/\lambda$ is real, extension to $\lambda \in \Lambda_{A,B}$ is straightforward.

We write $\rho = (x,y,\theta)$, $\phi(\rho) = -(\varphi(x,\theta) - y\theta)$. Stationary point of ϕ with respect to y,θ is given by $(y,\theta) = (0,0)$. If $M_0 u = v$ we have,

$$v(x) = h^{-n}\iint_{\Gamma(x)} e^{i(\phi_r(\rho)+\psi_r(\rho))/h}\left(b^r(\rho,\lambda) + c^r(\rho,\lambda)\right)u(y)dyd\theta, \tag{54}$$

with a good integration contour $\Gamma(x)$ passing through the critical point and $\psi_r(\rho) = \mathcal{O}(|\rho|^{2r+1})$, $c^r \sim \sum_{j=0}^\infty \lambda^{-j}c_j^r$ with, for $0 \le j \le r$, $c_j^r = \mathcal{O}(|\rho|^{2(r-j)+1})$. If $|\rho| \le \mathcal{O}(s^{1/2})$ we get

$$\psi_r(\rho) = \mathcal{O}(s^{r+1/2}),\ c^r(\rho) = \mathcal{O}(s^{r+1/2}).$$

Let $\Omega_2 \subset B(x_0, c_2 s^{1/2})$, be a neighborhood of $x_0 = 0$, and let

$$\Omega_1 = \pi_y\kappa^{-1}(\pi_x^{-1}\Omega_2 \cap \Lambda_\Phi) \subset\subset B(y_0, c_1 s^{1/2}),\ y_0 = 0.$$

By modifying $M_0 u$ with $\mathcal{O}(h^{N_0})$ in $L^2_\Phi(\Omega_2)$, with some $N_0 > 0$, we can restrict the integration contour $\Gamma(x)$ in (54) to $|y| + |\theta| \le \epsilon s^{1/2}$, with ϵ small enough.

It is clear that $e^{-i\lambda F^r(I;1/\lambda)}$ is well defined as a bounded operator $H_\Phi(\Omega_1) \mapsto H_\Phi(\Omega_2)$.

We want to estimate the difference $M_0 u - e^{-i\lambda F^r(I;1/\lambda)}u$ on Ω_2.

We make substitution $x = s^{1/2}\tilde{x}$, $y = s^{1/2}\tilde{y}$, $\theta = s^{1/2}\tilde{\theta}$ and write $u(y) = \tilde{u}(\tilde{y})$, $v(x) = \tilde{v}(\tilde{x})$, $\tilde{\rho} = (\tilde{x}, \tilde{y}, \tilde{\theta})$ and $s^{1/2}\tilde{\rho} = (s^{1/2}\tilde{x}, s^{1/2}\tilde{y}, s^{1/2}\tilde{\theta})$. Then, if $M_0 u = v$ we have

$$\tilde{v}(\tilde{x}) = \left(\frac{h}{s}\right)^{-n} \iint e^{\frac{i}{h}(\phi_r(s^{1/2}\tilde{\rho}) + \mathcal{O}(s^{r+1/2}))} \left(b^r(s^{1/2}\tilde{\rho}, \lambda) + \mathcal{O}(s^{r+1/2})\right) \tilde{u}(\tilde{y}) d\tilde{y} d\tilde{\theta}$$

Let $\tilde{\Gamma}$ be the image of $\Gamma(x)$ by substitution. We have then $\tilde{x} \in \tilde{\Omega}_2 \subset B(0, c_2)$, and for $(\tilde{y}, \tilde{\theta}) \in \tilde{\Gamma}$ we have $|\tilde{y}| + |\tilde{\theta}| \leq \epsilon$. We write $\phi_r^s := \frac{1}{s}\phi_r(s^{1/2}\rho)$. Then on $\tilde{\Gamma}$

$$-s^{-1}\Phi(s^{1/2}\tilde{x}) - \operatorname{Im}\phi_r^s + s^{-1}\Phi(s^{1/2}\tilde{x}) \sim -\left(|\tilde{x}|^2 + |\tilde{y}|^2 + |\tilde{\theta}|^2\right) + \mathcal{O}(s^{r+1/2}h^{-1}).$$

Then, if

$$v_r(\tilde{x}) = \left(\frac{h}{s}\right)^{-n} \int_{\tilde{\Gamma}} e^{\frac{s}{h}\phi_r^h} b_0^r(s^{1/2}\tilde{x}, s^{1/2}\tilde{y}, s^{1/2}\tilde{\theta}) \tilde{u}(\tilde{y}) d\tilde{y} d\tilde{\theta}, \tag{55}$$

we get

$$\|\tilde{v} - v_r\|_{\Phi, \tilde{\Omega}_2} \leq c(\ln(1/h))^n s^{r+1/2} h^{-1} \|\tilde{u}\|_{\Phi, \tilde{\Omega}_1 + B(0,\epsilon)} = \mathcal{O}(h^N)\|\tilde{u}\|_{\Phi, \tilde{\Omega}_1 + B(0,\epsilon)},$$

where $\|.\|_{\Phi,\Omega}$ denotes the norm in $H_\Phi(\Omega)$. Here $N > 0$ can be chosen arbitrary large by choosing r large enough.

∎

7. DEFORMATION OF THE ESCAPE FUNCTION IN A NEIGHBORHOOD OF $(0,0)$ AND FINAL DEFINITION OF THE SPACE.

We write $h = 1/\lambda_1$, $\lambda_1 = \operatorname{Re}\lambda$. Let W_0 be the neighborhood of 0 of the size $c_0\sqrt{h\ln(1/h)}$ as in (22) (with appropriate identifications of the domains). Let $W_1 \supset\supset W_0$ be the $\lambda-$independent domain as in (26).

Operators $M_0 = B_r M_1 B_r^{-1}$, $e^{-i\lambda F^r(I;1/\lambda)} = M_0 - R^r$ can be extended to the whole λ-independent neighborhood of 0. Let $U = B_r F T$, where T is the Bargman transform with a global choice of phase (33), F is the analytic Fourier integral operator quantizing the symplectic change of coordinates near $a_1 = 0$, B_r - the Birkhoff transform up to order r. On $W_1 \setminus W_0$ we can estimate the norm of R^r as in (50). Together with Lemma 6 it implies

Lemma 7. *For any $N \in \mathbb{N}$ and $\lambda \in \Lambda_{A,B}$ there is $r \in \mathbb{N}$ such that*

$$UMu = e^{-i\lambda 2d} e^{-i\lambda F^r(I;1/\lambda)} Uu + e^{-i\lambda 2d} R^r u, \text{ in } L^2_\Phi(V_1), \text{ and}$$
$$\|e^{-i\lambda 2d} R^r u\|_{L^2_\Phi(V_1)} \leq \mathcal{O}_r(h^N)\|u\|_{H(\Lambda_{tG})}, \text{ where } \pi(W_1) = V_1.$$

In Section 3 we constructed a global escape function G in $T^*(\partial\Omega_1)$, such that $G(\kappa(\rho)) - G(\rho) \geq 0$ on $T^*\partial(\Omega_1)$ and such that we have strict inequality, outside an arbitrary small neighborhood of $(a_1, 0) \in T^*\partial\Omega_1$.

Let Ω_{int} and W be a λ-independent neighborhoods of $(a_1, 0)$ (after usual identification of the domains), such that $W_0 \subset\subset W \subset\subset \Omega_{\text{int}} \subset\subset W_1$. We define a preliminary space $\mathcal{H}_{\text{pre}}$ associated to the Lagrangian space Λ_{tG} as follows

$$u \in \mathcal{H}_{\text{pre}} \Leftrightarrow \left\{ \begin{array}{c} T^{HS}u \in L^2_{tG}(T^*\partial\Omega_1 \setminus W) \\ Uu \in H_{\Phi_t}(\Omega_{\text{int}}),\ \kappa_U(\Lambda_{tG}) = \Lambda_{\Phi_t}. \end{array} \right.$$

In some small neighborhood of $(0,0)$,

$$W_{00} = \{\rho,\ |\rho| \leq c_{00}\sqrt{h\ln(1/h)}\} \subset\subset W_0, \tag{56}$$

we deform Λ_{Φ_t} in the following way.

We have $\kappa_U := \kappa_{B_r} \circ \kappa_F \circ \kappa_T :\ \Lambda_{tG} \mapsto \Lambda_{t\hat{G}}$. We denote again $\hat{G}$ by G. With G we associate an IR-lagrangian manifold which is essentially

$$\Lambda_{tG} = \{(x,\xi) = \exp{(itH_G)}(y,\eta);\ (y,\eta) \in T^*(\mathrm{neigh}(0))\}$$

for $0 < t \leq 1$, small.

Locally we have $\Lambda_+ = \{\xi = 0\}$, $\Lambda_- = \{x = 0\}$. For $x^2 + \xi^2 \leq \mathcal{O}(1)h\ln(1/h)$ we have $G \sim (x^2 - \xi^2)/2$, and Λ_{tG} is given by

$$\left\{ \begin{array}{ll} x_j & = (\cos t)y_j - i(\sin t)\eta_j \\ \xi_j & = -i(\sin t)y_j + (\cos t)\eta_j \end{array} \right. ,\ y_j, \eta_j \in \mathbb{R},\ 1 \leq j \leq n.$$

Then $\kappa_U(\Lambda_{tG}) = \Lambda_{\Phi_t} = \{\xi = \frac{2}{i}\frac{\partial \Phi_t}{\partial x}\}$, $\Phi_t = \frac{1}{2}(\cot t)(\operatorname{Im} x)^2 + \frac{1}{2}(\tan t)(\operatorname{Re} x)^2$.

Notice that $\Phi_{\pi/4} = |x|^2/2$ and that the corresponding IR-manifold is given by $\xi = -i\overline{x}$. Following [5] we look for a new IR-manifold Λ which coincides with $\Lambda_{\pi/4} = \Lambda_{\Phi_{\pi/4}}$ in some neighborhood of $(0,0)$ for $x^2 + \xi^2 \leq \mathcal{O}(1)h\ln(1/h)$ and with Λ_{Φ_t} outside.

First we notice that if $q_1(x)$ and $q_2(x)$ are strictly convex quadratic forms then we can find smooth strictly convex function $\phi(x)$, with $\phi(x) = q_1(x)$ near 0 and with $\phi(x) = q_2(x)$ outside V, where V is any given neighborhood of 0. Apply this with $V = \{x \in \mathbb{C}^n,\ |x| < K\}$ for some $K = \mathcal{O}(\sqrt{h\ln(1/h)})$, $q_1(x) = \Phi_{\pi/4}(x)$, $q_2(x) = \Phi_t(x)$.

Then replace $\phi(x)$ by $\phi_\sigma = \left(\frac{\sigma}{K}\right)^2 \phi(\frac{xK}{\sigma})$, $0 < \sigma \ll 1$, in order to decrease the neighborhood of 0, where $\phi \neq \Phi_t$, even further while keeping ϕ bounded in C^2. We have $\sigma \sim \sqrt{h\ln(1/h)}$ and the derivatives of higher order than 2 are diverging as $h \to 0$. Thus we get that the deformation ϕ_σ is strictly convex function which is close to the original function in C^2.

The deformation of Λ_{Φ_t} is denoted $\Lambda_{\tilde{\Phi}_t}$ and the new weight function satisfies

$$\tilde{\Phi}_t(x) = |x|^2/2,\ |x|^2 \leq \epsilon h\ln(1/h),\ \tilde{\Phi}_t(x) = \Phi_t,\ |x|^2 \geq c_{00}h\ln(1/h),$$

where $0 < \epsilon \ll c_{00}$ are some constants.

The final version of the space is given by

$$u \in \mathcal{H} \ \Leftrightarrow\ \left\{ \begin{array}{c} T^{HS}u \in L^2_{tG}(T^*\partial\Omega_1 \setminus W) \\ Uu \in H_{\tilde{\Phi}_t}(\Omega_{\mathrm{int}}). \end{array} \right.$$

8. Model problem.

In this section we derive expressions for the coefficients in the expansion in powers k^{-j} of the solution of equation (19).

We suppose that μ_j satisfy the non resonance condition (52). Thus all μ_j and $\nu_j = e^{\mu_j}$ are different. We also have $\mu_j > 0 \Rightarrow \nu_j > 1$.

For $\lambda \in \Lambda_{A,B}$, as in (18), consider the equation

$$2d\lambda + \lambda F^r\left(\frac{2\alpha + 1}{2i\lambda}; 1/\lambda\right) = 2\pi k,\ \alpha \in \mathbb{Z}^n_+,\ k \in \mathbb{Z}.$$

We omit r in the notations and proceed in increasing the order of generality.

A. Assume $F(\imath; 1/\lambda) = G(\imath, \mu) = \sum_1^n \mu_i \imath_i$, $\imath = x\xi$, is independent of λ and linear in $\imath$.

$$2d\lambda + \lambda G\left(\frac{1}{2i\lambda}(2\alpha_1 + 1), \ldots, \frac{1}{2i\lambda}(2\alpha_n + 1), \mu\right) = 2\pi k,\ k \in \mathbb{Z}\ \Rightarrow$$

$$2d\lambda + \frac{1}{2i}\sum_1^n \mu_i(2\alpha_i + 1) = 2\pi k\ \Rightarrow\ \lambda(k,\alpha) = k\frac{\pi}{d} + \frac{i}{4d}\sum_1^n \mu_i(2\alpha_i + 1),\ (k,\alpha) \in \mathbb{Z} \times \mathbb{N}^n.$$

We have

$$\operatorname{Re}\lambda = \frac{\pi k}{d},\ \operatorname{Im}\lambda = \frac{1}{4d}\sum_{j=1}^{n}\mu_j(2\alpha_j+1)\ \Rightarrow\ |\alpha| \sim \operatorname{Im}\lambda,$$

as $\operatorname{Re}\lambda \to \infty$, $|\alpha| = O(|\operatorname{Im}\lambda|)$, $|\alpha| = O(k^{1-\epsilon})$, $|\alpha| = O(\log|k|)$, $k \sim \operatorname{Re}\lambda$.

B. Assume that $F(\imath) = G(\imath,\mu) + H(\imath)$, polynomial of degree r, is independent of λ, where G is as above and $H(\imath) = O(\imath^2)$. We have

$$H\left(\frac{y}{\lambda}\right) = \sum_{j=2}^{r}\lambda^{-j}h_j(y),\ |1/\lambda| \to 0,\ |y| \le |\lambda|^{1-\epsilon}, \tag{57}$$

for some $\epsilon > 0$, where $|h_j(y)| = O(|y|^j)$, $h_j(y) = \sum_{|\alpha|=j}\frac{1}{\alpha!}\partial^\alpha H(0)y^\alpha$ is a homogeneous polynomial of degree j. Then we have

$$\frac{\lambda}{k}\left(1 + \frac{1}{4id\lambda}\sum_{i=1}^{n}\mu_i(2\alpha_i+1) + \frac{1}{2d}\sum_{j=2}^{r}\lambda^{-j}h_j(\alpha)\right) = \frac{\pi}{d}, \tag{58}$$

where $h_j(\alpha) = h_j\left(\frac{2\alpha+1}{2i}\right) = \mathcal{O}\left(|\alpha|^j\right)$ is homogeneous polynomial in α.

It is clear that equation (58) has solution in the form

$$\frac{\lambda}{k} = \frac{\pi}{d} + \frac{a_1}{k} + \frac{a_2}{k^2} + \ldots \in \Lambda_{A,B}. \tag{59}$$

Then (58) transforms to

$$\frac{a_1}{k} + \frac{a_2}{k^2} + \ldots + \frac{1}{4id}\sum_{i=1}^{n}\mu_i(2\alpha_i+1) + \frac{\lambda}{2dk}\sum_{j=2}^{r}\lambda^{-j}h_j(\alpha) = 0,$$

where the last term is of order $\mathcal{O}(|\alpha|^2/k^2)$.

Thus we get

$$a_1 = -\frac{1}{i4d}\sum_{j=1}^{n}\mu_j(2\alpha_j+1) = \mathcal{O}(|\alpha|) = \mathcal{O}(\ln k).$$

The next equation is

$$\begin{gathered}\frac{a_2}{k^2} + \frac{a_3}{k^3}\ldots + \frac{1}{2d}\left(\frac{h_2(\alpha)}{\lambda k} + \frac{h_3(\alpha)}{\lambda^2 k} + \ldots + \frac{h_r(\alpha)}{\lambda^{r-1}k}\right) = 0\ \Leftrightarrow \\ \left(\frac{\pi}{d}k + a_1 + \frac{a_2}{k} + \ldots\right)\left(\frac{a_2}{k^2} + \frac{a_3}{k^3} + \ldots\right) + \frac{1}{2d}\left(\frac{h_2(\alpha)}{k} + \frac{h_3(\alpha)}{\lambda k} + \ldots + \frac{h_r(\alpha)}{\lambda^{r-2}k}\right) = 0\end{gathered}$$

which implies

$$a_2 = -\frac{1}{2\pi}h_2\left(\frac{2\alpha+1}{2i}\right) = \mathcal{O}(|\alpha|^2). \tag{60}$$

With this choice of coefficients we get

$$\frac{\pi}{d}k\left(\frac{a_3}{k^3} + \frac{a_4}{k^4} + \ldots\right) + \left(a_1 + \frac{a_2}{k} + \ldots\right)\left(\frac{a_2}{k^2} + \frac{a_3}{k^3} + \ldots\right) + \frac{1}{2d}\left(\frac{h_3(\alpha)}{\lambda k} + \ldots + \frac{h_r(\alpha)}{\lambda^{r-2}k}\right) = 0.$$

Considering the coefficients for k^{-2} we get

$$\frac{\pi}{d}a_3 + a_1a_2 + \frac{h_3(\alpha)}{d\pi} = 0\ \Rightarrow\ a_3 = -\frac{d}{\pi}a_1a_2 - \frac{h_3(\alpha)}{\pi^2}.$$

We get

$$a_3 = -\frac{1}{i8\pi^2} \cdot \sum_{i=1}^{n} \mu_i(2\alpha_i+1) \cdot h_2\left(\frac{2\alpha+1}{2i}\right) - \frac{1}{\pi^2} h_3\left(\frac{2\alpha+1}{2i}\right).$$

It is clear that in this way we get all a_j, $j \leq r$ dependent only on $\{h_k\}_{k\leq j}$ and with this choice $\lambda_0/k = \pi/d + k^{-1}a_1 + k^{-2}a_2 + \ldots + k^{-r}a_r$ satisfies

$$\frac{\lambda_0}{k}\left(1 + \frac{1}{4id\lambda_0}\sum_{i=1}^{n}\mu_i(2\alpha_i+1) + \frac{1}{2d}\sum_{j=2}^{r}\lambda_0^{-j}h_j(\alpha)\right) - \frac{\pi}{d} = \mathcal{O}\left(\frac{|\alpha|^{r+1}}{k^{r+1}}\right). \tag{61}$$

C. We write $h = 1/\lambda$. In the general case $F^r(\imath;h) = F_0(\imath) + hF_1(\imath) + \ldots + h^rF_r(\imath) =: F_0(\imath) + K(\imath;h)$, with $F_0(\imath)$ as before, F_j polynomial in $\imath$ of degree $(r-j)$.

Then $F(0;h)/h = K(0;h)/h = F_1(0) + hF_2(0) + \ldots + h^{r-1}F_r$ and we can decompose

$$\begin{aligned}\frac{F(hy;h)}{h} =& \frac{K(0;h)}{h} + G(y,\mu) + \frac{H(hy)}{h} + \frac{K(hy;h)-K(0;h)}{h} = \\ =& \frac{K(0;h)}{h} + G(y,\mu) + \frac{H(hy)}{h} + \sum_{j=1}^{r-1} h^j k_j(y),\end{aligned}$$

where k_j is a polynomial of degree $\leq j$ and $k_j(0) = 0$. We get

$$F\left(\frac{y}{\lambda};1/\lambda\right) = F(0;1/\lambda) + \frac{1}{\lambda}G(y,\mu) + \sum_{j=2}^{r}\lambda^{-j}k_{j-1}(y).$$

Combining this with (57) where h_{j+1} is a homogeneous polynomial of degree $j+1$, we get

$$F\left(\frac{y}{\lambda};1/\lambda\right) = F(0;1/\lambda) + \frac{1}{\lambda}G(y,\mu) + \sum_{j=2}^{r}\lambda^{-j}f_j(y),$$

where for $2 \leq j \leq r$, $f_j(y) = h_j(y) + k_{j-1}(y)$ is a polynomial of degree $\leq j$. We have for example

$$f_2(y) = F_1'(0)y + \frac{1}{2}F_0''(0)y\cdot y.$$

We need to solve the equation

$$\frac{\lambda}{k}\left(1 + \frac{1}{4id\lambda}\sum_{i=1}^{n}\mu_i(2\alpha_i+1) + \frac{F_1(0)}{2d\lambda} + \frac{1}{2d}\sum_{j=2}^{r}\lambda^{-j}\left(F_j(0) + f_j(\alpha)\right)\right) = \frac{\pi}{d},$$

where we write $f_j(\alpha) = f_j((2\alpha+1)/2i) = \mathcal{O}(|\alpha|^j)$.

Then we can get all coefficients in the expansion $\frac{\lambda_0}{k} = \frac{\pi}{d} + \frac{a_1}{k} + \frac{a_2}{k^2} + \ldots + \frac{a_r}{k^r}$ such that

$$\frac{\lambda_0}{k}\left(1 + \frac{1}{4id\lambda_0}\sum_{i=1}^{n}\mu_i(2\alpha_i+1) + \frac{F_1(0)}{2d\lambda_0} + \frac{1}{2d}\sum_{j=2}^{r}\lambda_0^{-j}\left(F_j(0) + f_j(\alpha)\right)\right) - \frac{\pi}{d} = \mathcal{O}\left(\frac{|\alpha|^{r+1}}{k^{r+1}}\right).$$

First we get

$$a_1 = -\frac{1}{i4d}\sum_{i=1}^{n}\mu_i(2\alpha_i+1) - \frac{F_1(0)}{2d}.$$

Let $F_j(0)+f_j(\alpha)=q_j(\alpha)$ and fix $2\leq m\leq r$. Suppose that we have already chosen $a_1,\dots,a_{m-1}$ such that

$$\frac{\lambda_0}{k}\left(1+\frac{1}{4id\lambda_0}\sum_{i=1}^{n}\mu_i(2\alpha_i+1)+\frac{F_1(0)}{2d\lambda_0}+\frac{1}{2d}\sum_{j=2}^{r}\lambda_0^{-j}q_j(\alpha)-\frac{\pi}{d}\right)=\mathcal{O}\left(\frac{|\alpha|^m}{k^m}\right). \tag{62}$$

The left hand side of (62) is then equal to

$$\frac{a_2}{k^2}+\frac{a_3}{k^3}+\ldots+\frac{a_r}{k^r}+\frac{1}{2d}\left(\frac{q_2(\alpha)}{\lambda k}+\frac{q_3(\alpha)}{\lambda_0^2 k}+\ldots+\frac{q_r(\alpha)}{\lambda^{r-1}k}\right).$$

Denote

$$E_{m-1}(\lambda_0)=\frac{a_2}{k^2}+\frac{a_3}{k^3}+\ldots+\frac{a_{m-1}}{k^{m-1}}+\frac{1}{2d}\left(\frac{q_2(\alpha)}{\lambda_0 k}+\frac{q_3(\alpha)}{\lambda_0^2 k}+\ldots+\frac{q_{m-1}(\alpha)}{\lambda_0^{m-2}k}\right).$$

Then equation (62) writes

$$E_{m-1}(\lambda_0)+\sum_{j=m}^{r}\frac{a_j}{k^j}+\frac{1}{2d}\sum_{j=m}^{r}\frac{q_j(\alpha)}{\lambda_0^{j-1}k}=\mathcal{O}\left(\frac{|\alpha|^m}{k^m}\right). \tag{63}$$

Let $f(a_1,a_2,\dots,a_{m-1})$ be defined by

$$E_{m-1}(\lambda_0)\sim\frac{f(a_1,a_2,\dots,a_{m-1})}{k^m}+\mathcal{O}\left(\frac{|\alpha|^{m+1}}{k^{m+1}}\right)\ \text{as } |\alpha|/k\to 0.$$

Then equation (63) writes

$$\frac{f(a_1,a_2,\dots,a_{m-1})}{k^m}+\mathcal{O}\left(\frac{|\alpha|^{m+1}}{k^{m+1}}\right)+\frac{a_m}{k^m}+\mathcal{O}\left(\frac{|\alpha|^{m+1}}{k^{m+1}}\right)+\frac{1}{2d}\frac{q_m}{\lambda_0^{m-1}k}=-\sum_{j=m+1}^{r}\frac{q_j(\alpha)}{\lambda_0^{j-1}k}.$$

We chose a_m such that

$$f+a_m+\frac{1}{2d}\left(\frac{d}{\pi}\right)^{m-1}q_m(\alpha)=0\ \Leftrightarrow\ a_m=-f(a_1,a_2,\dots,a_{m-1})-\frac{1}{2d}\left(\frac{d}{\pi}\right)^{m-1}q_m(\alpha).$$

With this choice λ_0 satisfies equation (62) with the right hand side $\mathcal{O}\left(\frac{|\alpha|^{m+1}}{k^{m+1}}\right)$.

We summarize in the following theorem.

Theorem 5. *Let $\Lambda_{A,B}$ be defined in (18). For $r\in\mathbb{Z}_+$ let F^r be defined in (17). Then for any $(k,\alpha)\in\mathbb{N}\times\mathbb{N}^n$ such that $|\alpha|=\mathcal{O}(\ln k)$ and k large, there exist functions $a_j=a_j(\alpha)=\mathcal{O}(|\alpha|^j)$, polynomial in α, $j=0,1,\dots,r$, such that if we denote*

$$\lambda_r:=k\left(a_0+\frac{a_1}{k}+\frac{a_2}{k^2}+\ldots+\frac{a_{r+1}}{k^{r+1}}\right)\in\Lambda_{A,B},\ a_0=\frac{\pi}{d},\ a_1=-\frac{1}{i4d}\sum_{i=1}^{n}\mu_i(2\alpha_i+1)-\frac{F_1(0)}{2d},\dots, \tag{64}$$

then we have

$$2d\lambda_r+\lambda_r F^r\left(\frac{2\alpha+1}{2i\lambda_r};1/\lambda_r\right)-2\pi k=\mathcal{O}\left(\frac{(\ln k)^{r+2}}{k^{r+1}}\right)$$

and the coefficients a_j, are independent of r for $j=1,\dots,r+1$.

Note 1.

(1) *For any $r,k\in\mathbb{N}$ large and $\alpha=\mathcal{O}(\ln k)$, $\lambda_0:=\lambda_r(\alpha,k)$ satisfies*

$$1-e^{-i2d\lambda_0}K_\alpha^r(\lambda_0)=\mathcal{O}\left(\frac{(\ln k)^{r+2}}{k^{r+1}}\right),\ \textit{where } K_\alpha^r(\lambda):=e^{-i\lambda F^r\left(\frac{2\alpha+1}{2i\lambda};\lambda\right)}.$$

(2) *Using that operator M is microlocally unitary we know that $F_1(0)$ is real.*

(3) *For large k and $\alpha = \mathcal{O}(\ln k)$, we have* $\operatorname{Re}\lambda(\alpha, k+1) - \operatorname{Re}\lambda(\alpha, k) = \pi/2 + \mathcal{O}(1/k)$. *For any $\beta = \mathcal{O}(\ln k)$, $\beta \neq \alpha$, we have*

$$\operatorname{Im}\lambda(\alpha, k) - \operatorname{Im}\lambda(\beta, k) = \frac{1}{2d}\mu \cdot (\alpha - \beta) + \mathcal{O}\left(\frac{\ln^2 k}{k}\right),\ k >> 1.$$

In order to have good separation of strings as $k \to \infty$ (the first term in the above formula is dominated over the second error term) we need to impose the Diophantine condition:

$$\alpha \neq \beta,\ |\alpha|, |\beta| \leq m \ \Rightarrow\ |\mu\alpha - \mu\beta| \geq \frac{1}{C(D)} e^{-Dm},\ D > 0. \tag{65}$$

It would imply for any $\alpha \neq \beta$,

$$|\alpha|, |\beta| \leq c \ln k \ \Rightarrow\ |\mu\alpha - \mu\beta| \geq \frac{1}{C(D)} e^{-Dc\ln k} \geq \frac{1}{C(D)\, k^{cD}} >> \mathcal{O}\left(\frac{\ln^2 k}{k}\right),\ cD < 1.$$

As the next term is of order k^{-1} we have separation.

9. The first local Grushin problem in W_0.

9.1. **Notations.** In Ω_{int} for any $N \in \mathbb{N}$ we can chose $r \in \mathbb{N}$ such that we have

$$UMu = e^{-i\lambda 2d} e^{-i\lambda F^r(I;1/\lambda)} Uu + \mathcal{O}\left(h^N\right) |u|_{\mathcal{H}}.$$

Monomials x^α are formal eigenfunctions for $e^{-i\lambda F^r(I;1/\lambda)}$. Let K_α denote the corresponding eigenvalue: $e^{-i\lambda F^r(I;1/\lambda)} x^\alpha = K_\alpha x^\alpha$. We have

$$K_\alpha(\lambda) = e^{-i\lambda F^r(\frac{2\alpha_1+1}{2i\lambda}, \dots, \frac{2\alpha_n+1}{2i\lambda}; 1/\lambda)} =$$

$$= \exp\{-\frac{1}{2}\sum_{i=1}^{n} \mu_i(2\alpha_i + 1) - iF_1(0) - i\sum_{j=2}^{r} \lambda^{-j+1}\left(F_j(0) + f_j(\alpha)\right)\}, \tag{66}$$

(67) where $f_j(\alpha)$ is polynomial of degree $\leq j$.

We can have $K_\alpha(\lambda) = K_{\alpha'}(\lambda)$ for $\alpha \neq \alpha'$. We chose a value $\lambda_0 \in \Lambda_{A,B}$ and a multi-index α_0 such that

$$1 - e^{-i2d\lambda_0} K_{\alpha_0}(\lambda_0) = 0. \tag{68}$$

Let $\beta \in \mathbb{N}^n$ be such that $K_\beta(\lambda_0) \neq K_{\alpha_0}(\lambda_0)$.

We suppose that the Diophantine condition (65) in Note 1 in the previous section be satisfied and write $|e^{-i2d\lambda_0} K_\beta(\lambda_0)| = e^{|\mu(\alpha_0 - \beta)| + r}$. Then if $|\beta|,\ |\alpha_0| \leq C \ln \operatorname{Re}\lambda$ the Diophantine condition implies

$$|\mu(\alpha_0 - \beta)| \geq c(\operatorname{Re}\lambda)^{-\delta},\ r = o\left((\operatorname{Re}\lambda)^{-\delta}\right), \tag{69}$$

with some $0 < \delta < 1$. If $|\mu(\alpha_0 - \beta)| = \mathcal{O}\left((\operatorname{Re}\lambda)^{-\delta}\right)$ then we have

$$|1 - e^{-i2d\lambda_0} K_\beta(\lambda_0)| = \mathcal{O}\left((\operatorname{Re}\lambda)^{-\delta}\right),\ 0 < \delta < 1.$$

Let λ belong to a small neighborhood Ω_{λ_0} of λ_0.

We chose Ω_{λ_0} sufficiently small such that for all $\beta \in \mathbb{N}^n$ with $K_\beta(\lambda_0) \neq K_{\alpha_0}(\lambda_0)$ we have

$$\lambda \in \Omega_{\lambda_0} \cap \Lambda_{A,B},\ \left|1 - e^{-i2d\lambda} K_\beta(\lambda)\right| \geq \epsilon_0 \text{ and } \left|1 - e^{-i2d\lambda} K_{\alpha_0}(\lambda_0)\right| \leq \epsilon_0,$$

with

$$\epsilon_0 = \mathcal{O}\left((\operatorname{Re}\lambda)^{-\delta}\right),\ 0 < \delta < 1. \tag{70}$$

For $\lambda \in \Omega_{\lambda_0} \cap \Lambda_{A,B}$ we have

$$\operatorname{Im} \lambda = \frac{1}{4d}\mu(2\alpha_0 + 1) - \frac{\operatorname{Im} F_1(0)}{2d} + \mathcal{O}\left((\operatorname{Re}\lambda)^{-\delta}\right),$$

where we have $\operatorname{Im} F_1(0) = 0$ (see Note 1). Let

$$J = \{\alpha \in \mathbb{N}^n;\ |\alpha| \leq C \ln \operatorname{Re}\lambda_0,\ K_\alpha(\lambda_0) = K_{\alpha_0}(\lambda_0)\} \text{ and } a = \operatorname{Card} J.$$

Then we have

$$a \leq \mathcal{O}(1) \ln \operatorname{Re}\lambda_0. \tag{71}$$

We can chose $\Lambda_{A,B}(\epsilon_0) \subset \Omega_{\lambda_0} \cap \Lambda_{A,B}$ such that

$$\text{for } \lambda \in \Lambda_{A,B}(\epsilon_0),\ \alpha \in J,\ \left|1 - e^{-i2d\lambda} K_\alpha(\lambda)\right| \leq 2\epsilon_0. \tag{72}$$

We will use the following notations: $\lambda_1 = \operatorname{Re}\lambda$, $h = 1/\lambda_1$ and $s = h\ln(1/h)$.

9.2. Monomials form an almost orthonormal base in H_Φ.

For $\Omega \subset\subset \mathbb{C}^n$ we denote

$$H_\Phi(\Omega) = \{u \in \mathcal{H}ol(\Omega) \text{ such that } \int_\Omega |u|^2 e^{-2\Phi/h} L(dx) < \infty\},$$

where Φ is the modified weight defined in Section 7 such that $\Phi = |x|^2/2$ for $x \in B(0, \epsilon\sqrt{h\ln(1/h)})$, $\epsilon < c_{00}$.

Since the weight function Φ is fixed from now on, we shall suppress it from our notations as much as possible. Normalized monomials

$$\varphi_\alpha(x) = c_\alpha h^{-n/2}(h^{-1/2}x)^\alpha,\ c_\alpha := (\pi^n \alpha!)^{-1/2}$$

form an orthonormal base in $H_{\Phi_0}(\mathbb{C}^n)$ for $\Phi_0 = |x|^2/2$.

Following [5] we show that $\varphi_\alpha(x)$ are orthonormalized in $H_\Phi(\Omega)$ with an error of order h^N for any $N \in \mathbb{N}$.

Lemma 8. *For all $\Omega \subset B(0,c)$, where c is some real number, we have*

$$\langle \varphi_\alpha | \varphi_\beta \rangle_{H_\Phi(\Omega)} = \delta_{\alpha,\beta} + \mathcal{O}(h^N)$$

for $N > 0$, which can be taken arbitrary large by choosing c_{00} large enough, and $\mathcal{O}$ is uniform in α, β, Ω.

Proof:

$$\int_\Omega \left(\Pi_{j=1}^n |x_j|^{2\alpha_j}\right) e^{-2\Phi(x)/h} L(dx) = \Pi_{j=1}^n \int_{\mathbb{C}} |x_j|^{2\alpha_j} e^{-x_j^2/h} L(dx_j) -$$
$$- \int_{|x| \geq \epsilon\sqrt{h\ln(1/h)}} \left(\Pi_{j=1}^n |x_j|^{2\alpha_j}\right) e^{-|x|^2/h} L(dx) + \int_{\{x\in\Omega,\ |x| \geq \epsilon\sqrt{h\ln(1/h)}\}} \left(\Pi_{j=1}^n |x_j|^{2\alpha_j}\right) e^{-2\Phi(x)/h} L(dx) =$$
$$= I_1 - I_2 + I_3$$

and ϵ is such that $\phi(x) = |x|^2/2$ for $|x| \leq \epsilon\sqrt{h\ln(1/h)}$. The first term I_1 is equal to $\pi^n \alpha! h^{|\alpha|+n}$. We need to show that

$$\frac{|I_2| + |I_3|}{\pi^n \alpha! h^{|\alpha|+n}} = \mathcal{O}(h^N)$$

for $h \to 0$ uniformly in α. As $\Phi \geq c|x|^2$ for some $c > 0$, it is enough to show the following:

$$c_\alpha^2 \int_{\{x \in \mathbb{C}^n,\ |x| \geq \epsilon\sqrt{h\ln(1/h)}\}} \left(h^{-1/2}x\right)^{2\alpha} e^{-2c|x|^2/h} L(dx) = c_\alpha^2 \left(\frac{h}{c}\right)^n c^{-|\alpha|} \int_{y \geq \epsilon\sqrt{c\ln(1/h)}} y^{2\alpha} e^{-2|y|^2} L(dy)$$

$$\leq c_\alpha^2 \left(\frac{h}{c}\right)^n c^{-|\alpha|} e^{-\epsilon^2 c \ln(1/h)} \underbrace{\int_{\mathbb{C}^n} e^{-|y|^2} y^{2\alpha} L(dy)}_{\pi^n \alpha! \equiv c_\alpha^{-2}} = \mathcal{O}(h^N),$$

where we mean $x^\alpha = \Pi_{j=1}^n x_j^{\alpha_j}$ and make change of variables $y = \sqrt{\frac{c}{h}}x$.

Next we show approximate orthogonality:

$$\langle \varphi_\alpha | \varphi_\beta \rangle_{H_\Phi(\Omega)} = \int_{|x| \leq \epsilon\sqrt{h\ln(1/h)}} \varphi_\alpha \overline{\varphi}_\beta e^{-|x|^2/h} L(dx) +$$

$$+ \int_{x \in \Omega,\ |x| \geq \epsilon\sqrt{h\ln(1/h)}} \varphi_\alpha \overline{\varphi}_\beta e^{-2\Phi(x)/h} L(dx) = I_1 + I_2,$$

$I_1 = 0$, $I_2 = \mathcal{O}(h^N)$ by previous calculus. ∎

9.3. **Grushin problem.** We denote $V_0 = \pi(W_0)$, where W_0 is the image by κ_U of the domain defined in (22).

For $N \in \mathbb{N}$, we introduce the expansions:

$$\tau_N u(x) = \sum_{|\alpha| < N} (\alpha!)^{-1} (\partial_x^\alpha u)(0) x^\alpha = \sum_{|\alpha| < N} (\alpha!)^{-1} ((h^{1/2}\partial_x)^\alpha u)(0) (h^{-1/2}x)^\alpha.$$

We denote $F^N := \mathcal{I}m\, \tau_N$, $O^N := \mathcal{K}er\, \tau_N$, $M(N) = \dim(F^N)$.

Suppose that $N \in \mathbb{N}$ satisfy

$$\max_{\alpha \in J} |\alpha| \leq N = \mathcal{O}(1) \ln \operatorname{Re} \lambda_0, \tag{73}$$

where λ_0 is fixed as before satisfying (68).

In addition, we need the expansions

$$\tau_J u(x) = \sum_{\alpha \in J} (\alpha!)^{-1} (\partial_x^\alpha u)(0) x^\alpha = \sum_{\alpha \in J} (\alpha!)^{-1} ((h^{1/2}\partial_x)^\alpha u)(0) (h^{-1/2}x)^\alpha.$$

We denote $F_J := \mathcal{I}m\, \tau_J$, $O_J := \mathcal{K}er\, \tau_J$. We have $a(\epsilon_0) = \dim(F_J)$.

We have introduced the approximately orthonormal basis of monomials:

$$\varphi_\alpha(x) = c_\alpha h^{-n/2} (h^{-1/2}x)^\alpha,\ c_\alpha = (\pi^n \alpha!)^{-1/2}.$$

Then

$$(\alpha!)^{-1}((h^{1/2}\partial_x)^\alpha u)(0)(h^{-1/2}x)^\alpha = \varphi_\alpha \cdot (\pi h)^{n/2} (\alpha!)^{-1/2} ((h^{1/2}\partial_x)^\alpha u)(0).$$

Let $R_- :\ \mathbb{C}^a \mapsto F_J,\ R_+ :\ F_J \mapsto \mathbb{C}^a$ be defined by the formulas

$$u \in F_J,\ u_- = \{u_{-,\alpha}\}_{|\alpha \in J} \in \mathbb{C}^a,\ R_- u_- = \sum_{\alpha \in J} u_{-,\alpha} \varphi_\alpha,$$

$$R_+ u = \left\{(\pi h)^{n/2} (\alpha!)^{-1/2} (h^{1/2}\partial_x)^\alpha u(0)\right\}_{\alpha \in J}.$$

By direct calculation we get $R_+ u = \langle u, \varphi_\alpha \rangle_{\alpha \in J} (1 + \mathcal{O}(h^N))$.

Let

$$\|u\|_\Omega := \|u\|_{H_\Phi(\Omega)},\ |x|_{\mathbb{C}^a} := \sqrt{\sum_{\alpha \in J} |x_\alpha|^2}.$$

We write $\|.\|$ for natural operator norm. We have

$$R_+R_- = Id_{|\mathbb{C}^a},\ R_-R_+ = \tau_J = Id_{|F_J},\ \tau_N R_- = R_-,\ R_+\tau_N = R_+,$$
$$\|R_-\| = 1 + \mathcal{O}(h^N),\ \|R_+\| = 1 + \mathcal{O}(h^N).$$

The estimates on the norms of operators $R_\pm$ follow from Lemma 8.

We pose the first Grushin problem for $\lambda \in \Lambda_{A,B}(\epsilon_0)$:

$$\left\{ \begin{array}{l} \tau_N(I - e^{i2d\lambda}e^{-i\lambda F^r(I;1/\lambda)})\tau_N u + R_- u_- = v, \\ R_+ u = v_+,\ u, v \in F^N. \end{array} \right. \tag{74}$$

Theorem 6. *For a given* $(v, v_+) \in F^N \times \mathbb{C}^a$ *and all* $\lambda \in \Lambda_{A,B}(\epsilon_0)$, *the Grushin problem (74) has unique exact solution* $(u, u_-) \in F^N \times \mathbb{C}^a$, *which satisfies*

$$\|u\|_{H_\Phi(\Omega_1)} + |u_-|_{\mathbb{C}^a} \leq \mathcal{O}\left((\operatorname{Re}\lambda)^\delta\right)(\|v\|_{H_\Phi(\Omega_2)} + |v_+|_{\mathbb{C}^a}),\ 0 < \delta < 1, \tag{75}$$

where δ *is as in (69),* $\Omega_2 \subset\subset \Omega_1 \subset\subset V_0$ *and* $|x|_{\mathbb{C}^a} := \sqrt{\sum_{\alpha\in J}|x_\alpha|^2}$.

Proof: We denote $h = 1/\operatorname{Re}\lambda$, $\lambda \in \Lambda_{A,B}(\epsilon_0)$, $z = e^{i2d\lambda}$. The solution operator for (74) is

$$\mathcal{E}^0 = \begin{pmatrix} E^0 & E^0_+ \\ E^0_- & E^0_{+-} \end{pmatrix},$$

where

$$\begin{aligned} E^0(v) &= \sum_{\alpha\notin J, |\alpha|<N} (1 - zK_\alpha)^{-1}(\alpha!)^{-1}((h^{1/2}\partial_x)^\alpha v)(0)(h^{-1/2}x)^\alpha \\ &= \sum_{\alpha\notin J, |\alpha|<N} (1 - zK_\alpha)^{-1}\langle v, \varphi_\alpha\rangle\varphi_\alpha\left(1 + \mathcal{O}(h^N)\right), \\ E^0_+(v_+) &= \sum_{\alpha\in J} v_{+,\alpha}(\pi^n\alpha!)^{-1/2}h^{-n/2}(h^{-1/2}x)^\alpha = \sum_{\alpha\in J} v_{+,\alpha}\varphi_\alpha(x) = R_-v_+, \\ E^0_-(v) &= \left((\pi h)^{n/2}(\alpha!)^{-1/2}(h^{1/2}\partial_x)^\alpha v(0)\right)_{\alpha\in J} = \langle v, \varphi_\alpha\rangle_{|\alpha\in J}\left(1 + \mathcal{O}(h^N)\right) = R_+v, \\ E^0_{+-}(v_+) &= -\{(1 - zK_\alpha(\lambda))\}_{|\alpha\in J}v_+, \end{aligned}$$

where $\{(1 - zK_\alpha(\lambda))\}_{|\alpha\in J}$ is diagonal matrix with entrees $(1 - zK_{\alpha_1}), \ldots, (1 - zK_{\alpha_N})$. Some of entrees can coincide. Solutions to (74) are $u = E^0(v) + E^0_+(v_+)$, $u_- = E^0_-(v) + E^0_{+-}(v_+)$.

Let $B_{\epsilon\sqrt{s}} \subset\subset \Omega_2 \subset\subset \Omega_1 \subset\subset V_0$, $s = h\ln(1/h)$. We have the following estimates:

$$\begin{aligned} \|E^0(v)\|^2_{\Omega_1} &\leq \epsilon_0^{-2} \sum_{\alpha\notin J, |\alpha|<N} (\pi h)^n(\alpha!)^{-1}\left|((h^{1/2}\partial_x)^\alpha v)(0)\right|^2(1 + \mathcal{O}(h^N)) \leq \\ &\leq \epsilon_0^{-2}\|v\|^2_{\Omega_2}(1 + \mathcal{O}(h^N)),\ \|E^0_+(v_+)\|^2_{\Omega_1} = \sum_{\alpha\in J}|v_{+,\alpha}|^2(1 + \mathcal{O}(h^N)), \end{aligned} \tag{76}$$

as $E^0_+(v_+) = R_-v_+$.

We get

$$\|u\|_{\Omega_1} = \|E^0(v) + E^0_+(v_+)\|_{\Omega_1} \leq \epsilon_0^{-1}\left(1 + \mathcal{O}(h^N)\right)(\|v\|_{\Omega_2} + |v_+|_{\mathbb{C}^a}). \tag{77}$$

As $E^0_-(v) = R_+v$, we have already proven that

$$|E^0_-(v)|^2_{\mathbb{C}^a} \leq \left(1 + \mathcal{O}(h^N)\right)\|v\|^2_{\Omega_2}.$$

By (71), (72) and (70) we have also

$$|E^0_{+-}(v_+)|_{\mathbb{C}^a} \leq Ca\epsilon_0|v_+|_{\mathbb{C}^a} \leq \mathcal{O}(1)\ln\operatorname{Re}\lambda\cdot(\operatorname{Re}\lambda)^{-\delta}|v_+|_{\mathbb{C}^a},\ 0 < \delta < 1.$$

It implies

$$|u_-|_{\mathbb{C}^a} = |E_-^0(v)|_{\mathbb{C}^a} + |E_{+-}^0(v_+)|_{\mathbb{C}^a} \le (1+\mathcal{O}(h^N))(\|v\|_{\Omega_2} + |v_+|_{\mathbb{C}^a}). \tag{78}$$

Adding inequalities (77) and (78) and using (70) we get the estimate. ∎

As the Grushin problem (74) is invertible then it is well posed. Let

$$P_N(\lambda) := \tau_N(I - e^{i2d\lambda} e^{-i\lambda F^r(I;1/\lambda)}(\lambda))\tau_N.$$

By general consideration it follows that $E_+^0 : \mathcal{K}er\, E_{-+}^0 \mapsto \mathcal{K}er\, P_N(\lambda)$ is a bijection. As $P_N(\lambda)$ is a family of Fredholm operators depending holomorphically on $\lambda \in \Lambda_{A,B}(\epsilon_0)$ then

$$\Lambda_{A,B}(\epsilon_0) \ni \lambda \ \mapsto \ (P_N(\lambda))^{-1}$$

is a meromorphic family of operators.

We have the formula

$$(P_N(\lambda))^{-1} = E^0(\lambda) - E_+^0(\lambda)\left(E_{-+}^0(\lambda)\right)^{-1} E_-(\lambda).$$

Let γ be simple loop such that $E_{-+}^0(\lambda)$ is invertible for $\lambda \in \gamma$.

The number of poles $n(\gamma)$ of $P_N(\lambda)^{-1}$ inside γ counted with there multiplicity is given by

$$n(\gamma) = \frac{1}{2\pi i}\operatorname{tr}\int_\gamma (P_N(\lambda))^{-1}\,d\lambda = \frac{1}{2\pi i}\int_\gamma \operatorname{tr}\left(E_+^0\left(E_{-+}^0\right)^{-1}E_-\right)d\lambda.$$

From the other side, the number of roots of $\det E_{-+}^0$ inside γ is equal to

$$m(\gamma) = \frac{1}{2\pi i}\int_\gamma \frac{\left(\det E_{-+}^0(\lambda)\right)'}{\det E_{-+}^0}\,d\lambda = \frac{1}{2\pi i}\int_\gamma \operatorname{tr}\left\{\left(E_{-+}^0\right)'\left(E_{-+}^0\right)^{-1}\right\}d\lambda.$$

As in [8] we have then $n(\gamma) = m(\gamma)$.

10. Some useful estimates.

10.1. Estimate on $u_N \in \mathcal{K}er\,\tau_N$.

Let $s = h\ln(1/h)$, $h = 1/\lambda_1$, $\lambda_1 = \operatorname{Re}\lambda$, $\lambda \in \Lambda_{A,B}$, $B_r = B(0,r) = \{x,\ |x| \le r\}$.

Lemma 9. *Let $N \in \mathbb{N}$, $N = \mathcal{O}(\ln \operatorname{Re}\lambda)$, satisfying (73). Let $u_N \in H_\Phi(B_{\epsilon\sqrt{s}}) \cap \mathcal{K}er\,\tau_N$. Put*

$$r_m := \frac{\sqrt{N+1/2}}{2\sqrt{s\lambda_1}} = \mathcal{O}(1).$$

Let $\epsilon \ge r_m$ be such that $\Phi = |x|^2/2$ for $x \in B_{\epsilon\sqrt{s}}$. Then for any $r_0 \le r_m$,

$$\|u_N\|_{B_{r_0\sqrt{s}}} \le \|u_N\|_{B_{\epsilon\sqrt{s}} \setminus B_{r_0\sqrt{s}}}.$$

Proof: We need to prove

$$\int_{B_{r_0\sqrt{s}}} e^{-|x|^2/h}|u_N(x)|^2 L(dx) \le \int_{\complement B_{r_0\sqrt{s}} \cap B_{\epsilon\sqrt{s}}} e^{-|x|^2/h}|u_N(x)|^2 L(dx).$$

We change the variables $x = \tilde{x}\sqrt{s}$ in order to have the domain of integration independent of h :

$$x \in B_{\epsilon\sqrt{s}} \ \Leftrightarrow\ \tilde{x} \in B_\epsilon.$$

Denote $\Phi_s = \frac{1}{s}\Phi(\sqrt{s}x)$ and

$$H_{\Phi_s}(B_\epsilon) = \mathcal{H}ol\,(B_\epsilon) \cap L_{\Phi_s}(B_\epsilon) = \{v \in \mathcal{H}ol\,(B_\epsilon),\ \int_{B_\epsilon} |v|^2 e^{-s2\Phi_s/h} L(dx) < \infty\}.$$

Let $u \in H_\Phi(B_\epsilon)$, $u_N(x) = u(x) - \sum_{|\alpha|<N} \frac{\partial_x^\alpha u(0)}{\alpha!} x^\alpha$, $B_\epsilon := B(0,\epsilon)$.

Suppose that $\epsilon > 0$ is small enough such that $\Phi = |x|^2/2$ on the domain of integration. Thus $\Phi_s = |x|^2/2$. We must prove that for $(s\lambda_1)^{-1/2}\sqrt{N+1/2} \le \epsilon$ and $r_m = 1/2(s\lambda_1)^{-1/2}\sqrt{N+1/2}$ we have

$$\int_{B_{r_m}} e^{-s\lambda_1|x|^2}|u_N(x)|^2 L(dx) \le \int_{\complement B_{r_m}\cap B_\epsilon} e^{-s\lambda_1|x|^2}|u_N(x)|^2 L(dx).$$

The monomials constitute an orthogonal base in $H_\Phi(B_\epsilon)$ and it is enough to show the inequality for x^α, $|\alpha| \ge N$. Let $\alpha_j \in \mathbb{N}$ be such that $\alpha_j \ge N$. For $u_N(x) = x_j^{\alpha_j}$ we have

$$\int_{|x|\le\epsilon} e^{-s\lambda_1|x|^2}|f_N(x)|^2 L(dx) = C_n(s\lambda_1)^{-n/2}\int_{r\le\epsilon\sqrt{s\lambda_1}} e^{-r^2}r^{2\alpha_j+1}dr.$$

As

$$\left(e^{-r^2}r^{2\alpha_j+1}\right)'_r = (-2r^2+2\alpha_j+1)e^{-r^2}r^{2\alpha_j} = 0 \iff r = r_m(\alpha) := \sqrt{\alpha_j+1/2}$$

we get that the function $e^{-r^2}r^{2\alpha+1}$ is increasing for $0 \le r \le r_m$. Then

$$\int_0^{r_m/2} e^{-r^2}r^{2\alpha_j+1}dr \le \int_0^{r_m/2} e^{-(r+r_m/2)^2}\left(r+\frac{r_m}{2}\right)^{2\alpha_j+1}dr = \int_{r_m/2}^{r_m} e^{-r^2}r^{2\alpha_j+1}dr.$$

For all α such that $|\alpha| > N$ the maximum $r_m(\alpha) > r_m(\alpha_N)$ with $|\alpha_N| = N$. Thus, if the above estimate is valid for $|\alpha_N| = N$ then it implies that it is valid for any $|\alpha| > N$. ■

10.2. **Model estimate.**

Lemma 10. *As before we write $s = h\ln(1/h)$, $h = 1/\lambda_1$, $\lambda_1 = \operatorname{Re}\lambda$. Let Ω be any λ-independent neighborhood of 0 and $r_0 > 0$ such that $B_{r_0\sqrt{s}} \subset\subset \Omega$.*

Let $A = \operatorname{diag}(e^{\mu_1},\dots,e^{\mu_n})$ be as in Theorem 3. Denote $A^{-1}(\Omega) := \{A^{-1}x,\ x\in\Omega\}$.

For any $c_0 > 0$ and $|z| \le e^{c_0\ln\lambda_1}$ and r_0 large enough there exists $m \le 1/2$ such that

$$\int_{\Omega\setminus B_{r_0\sqrt{s}}} e^{-2\Phi(x)/h}|ze^{-\sum_{j=1}^n \mu_j(x_j\partial_j+1/2)}u|^2 L(dx) \le m^2\int_{A^{-1}(\Omega)\setminus A^{-1}(B_{r_0\sqrt{s}})} e^{-2\Phi(x)/h}|u|^2 L(dx).$$

Proof: We have

$$e^{-\sum_{j=1}^n \mu_j(x_j\partial_j+1/2)}u = e^{-\sum_{j=1}^n \mu_j/2}u(A^{-1}x),\ A = \operatorname{diag}(e^{\mu_1}\dots e^{\mu_n}).$$

Let first $\Phi = |x|^2/2$. We put for simplicity $n = 1$, $\mu_j = \mu$. Then $e^{-\mu(x\partial+1/2)}u = e^{-\mu/2}u(e^{-\mu}x)$. The general case is straightforward. We change the variables $\tilde{x} = e^{-\mu}x$, $L(dx) = e^{2\mu}L(d\tilde{x})$, and denote the new variable $\tilde{x}$ again by x. Let $A^{-1}(\Omega) := \{e^{-\mu}x,\ x\in\Omega\}$. Then we have, with $s = h\ln\lambda_1$,

$$e^{-\mu}\cdot\int_{\Omega\setminus B_{r_0\sqrt{s}}} e^{-|x|^2/h}|zu(e^{-\mu}x)|^2 L(dx) \le e^{2c_0\ln\lambda_1+\mu}\int_{A^{-1}(\Omega)\setminus B_{r_0\sqrt{s}e^{-\mu}}} e^{-e^{2\mu}|x|^2/h}|u(x)|^2 L(dx) =$$

$$= e^{2c_0\ln\lambda_1+\mu}\int_{A^{-1}(\Omega)\setminus B_{r_0\sqrt{s}e^{-\mu}}} e^{-(e^{2\mu}-1)|x|^2/h}e^{-|x|^2/h}|u(x)|^2 L(dx) \le$$

(79)

$$\le e^{2c_0\ln\lambda_1+\mu}e^{-(e^{2\mu}-1)e^{-2\mu}r_0^2 s/h}\int_{A^{-1}(\Omega)\setminus B_{r_0\sqrt{s}e^{-\mu}}} e^{-|x|^2/h}|u(x)|^2 L(dx) =$$

$$= m^2\int_{A^{-1}(\Omega)\setminus B_{r_0e^{-\mu}\sqrt{s}}} e^{-|x|^2/h}|u(x)|^2 L(dx)$$

with

$$m^2 := e^{2c_0 \ln \lambda_1 + \mu - \ln \lambda_1 (1-e^{-2\mu}) r_0^2}.$$

We have $m \leq 1/2$ if

$$e^{2c_0 \ln \lambda_1 + \mu - \ln \lambda_1 (1-e^{-2\mu}) r_0^2} \leq 2^{-2} \ \Leftrightarrow\ 2c_0 \ln \lambda_1 + \mu - \ln \lambda_1 (1-e^{-2\mu}) r_0^2 \leq -2 \ln 2 \ \Leftrightarrow$$

$$r_0 \geq \sqrt{\frac{2c_0 \ln \lambda_1 + \mu + 2 \ln 2}{\ln \lambda_1 \cdot (1-e^{-2\mu})}}.$$

For general weights Φ we use the strict convexity and get (after change of variables $\tilde{x} = e^{-\mu} x$)

$$\Phi(e^{\mu}\tilde{x}) - \Phi(\tilde{x}) \geq c|\tilde{x}|^2,\ c > 0,\ x \in V_1,$$

and in the above formula (79) exchange $e^{2\mu} - 1$ by c. ∎

10.3. **General estimate.** In this section we will write $h = 1/\lambda$ and suppose that λ is real. Generalization to $\lambda \in \Lambda_{A,B}$ is straightforward.

From (57) it follows that

$$e^{-iH^r(hy)/h} \sim 1 + hp_1(y) + h^2 p_2(y) + \dots,\ |y| \leq |h|^{\delta - 1/2},\ \delta > 0,$$

where the polynomial $p_j(y) = \sum_{|\alpha| \in [j+1, 2j]} q_{\alpha, j} y^{\alpha}$ is a linear combination of monomials of degree in $[j+1, 2j]$. Moreover, $p_1(y) = -\frac{i}{2} H''(0) y \cdot y$. Then

$$e^{-iF(hy)/h} = e^{-i\mu y} \left(1 + hp_1(y) + h^2 p_2(y) + \dots\right) = \left(1 + hp_1(i\partial_\mu) + h^2 p_2(i\partial_\mu) + \dots\right) e^{-i\mu y}.$$

Taking $y = h^{-1} \cdot I = \left(x\partial_x + 1/2\right)/i$, we have the following representation:

$$e^{-iF_0^r(I)/h} = e^{-iH^r(ih\partial_\mu)/h} e^{-\sum_{i=1}^n \mu_i (x_i \partial_i + 1/2)},\ F_0(\imath) = G(\imath, \mu) + H^r(\imath) = \sum_{i=1}^n \mu_i \cdot \imath_i + \mathcal{O}_r(\imath^2),$$

polynomial of degree r independent of h.

Then,

$$e^{-iH^r(ih\partial_\mu)/h} e^{-\sum_{i=1}^n \mu_i (x_i \partial_i + 1/2)} u = e^{-iH(ih\partial_\mu)/h} e^{-\sum_{j=1}^n \mu_j / 2} u(A^{-1}x),\ A = \operatorname{diag}(e^{\mu_1} \dots e^{\mu_n}).$$

For simplicity we put $n = 1$. We have

$$\begin{aligned} e^{-iF_0^r(I)/h} u &= e^{-iH(ih\partial_\mu)/h} e^{-\mu(x\partial + 1/2)} u(x) = \\ &= e^{-\mu/2} u(e^{-\mu}x) + h \sum_{j=1}^{r-1} h^{j-1} \sum_{l \in [j+1, 2j]} q_{l,j} \partial_\mu^l (e^{-\mu/2} u(e^{-\mu}x)) = \\ &= e^{-\mu/2} u(e^{-\mu}x) + e^{-\mu/2} T_0^r (e^{-\mu}x), \end{aligned}$$

$$\text{where } T_0^r u(x) := h \sum_{j=1}^{r-1} h^{j-1} \sum_{l \in [j+1, 2j]} q_{l,j} \sum_{k=0}^{l} c_{k,l} u^{(k)}(x) \cdot x^k = \mathcal{O}(h).$$

In general case

$$e^{-iF(hy;h)/h} = e^{-iK(0;h)/h} e^{-iG(y,\mu)} (1 + hq_1(y) + h^2 q_2(y) + \dots),$$

where $q_j(y)$ is a polynomial of degree at most $2j$, and as before, this can be written

$$e^{-iF(hy;h)/h} = e^{-iK(0;h)/h} (1 + hq_1(i\partial_\mu) + h^2 q_2(i\partial_\mu) + \dots) e^{-iG(y,\mu)}.$$

We will write it in the form

$$\frac{F(hy;h)}{h} = \frac{F(0;h)}{h} + G(y,\mu) + \frac{J(hy;h)}{h}$$

and then (see also [14], page 347)

$$e^{-iF(I(h);h)/h} = e^{-iF(0;h)/h} e^{-iJ(ih\partial_\mu;h)/h} e^{-\sum_{i=1}^{n}\mu_i(x_i\partial_i+1/2)}$$

and the difference from the case $F = G(I,\mu) + H(I)$ is only the factor $e^{-iF(0;h)/h}$, which is bounded as $h \to 0$, and that $q_j(y)$ is a polynomial of degree at most $2j$ and not just in $[j+1, 2j]$.

We have

$$\begin{aligned} & e^{-iF(I(h);h)/h}u = e^{-iF(0;h)/h} e^{-iJ(ih\partial_\mu;h)/h} e^{-\sum_{i=1}^{n}\mu_i(x_i\partial_i+1/2)}u = \\ & e^{-iF(0;h)/h}\left(e^{-\mu/2}u(e^{-\mu}x) + h\sum_{j=1}^{r-1} h^{j-1}\sum_{l=0}^{2j} q_{l,j}\partial_\mu^l(e^{-\mu/2}u(e^{-\mu}x))\right) = \\ & = e^{-iF(0;h)/h}\left(e^{-\mu/2}u(e^{-\mu}x) + e^{-\mu/2}T^r(e^{-\mu}x)\right), \qquad (80) \\ & \text{where } T^r u(x) := h\sum_{j=1}^{r-1} h^{j-1}\sum_{l=0}^{2j} q_{l,j}\sum_{k=0}^{l} c_{k,l}u^{(k)}(x)\cdot x^k. \end{aligned}$$

We put $z = e^{-i2d\lambda}$. Then for $\lambda \in \Lambda_{A,B}$ we have $|z| < e^{2dA\ln\operatorname{Re}\lambda}$, with A as in the definition of $\Lambda_{A,B}$.

We can apply Lemma 10 with $c_0 = 2dA$. Then, for r_0 large enough we have

$$\begin{aligned} & e^{-\mu}\cdot\int_{\Omega\setminus B_{r_0\sqrt{s}}} e^{-2\Phi(x)/h}|z|^2\left|u(e^{-\mu}x) + T^r u(e^{-\mu}x)\right|^2 L(dx)) \leq \\ & m^2\int_{A^{-1}(\Omega)\setminus A^{-1}(B_{r_0\sqrt{s}})} e^{-2\Phi(x)/h}\left|u(x) + T^r u(x)\right|^2 L(dx). \qquad (81) \end{aligned}$$

For any $\Omega_2 \subset\subset \Omega_1 \subset\subset \Omega$, we have for any $u \in H_\Phi(\Omega)$,

$$\|(h^{-1/2}x)^\alpha(h^{1/2}D_x)^\beta u\|_{\Omega_2} \leq \|\left(1 + h^{-1/2}|x|\right)^{|\alpha|+|\beta|}u\|_{\Omega_1},$$

see Corollary 4.2 in [8].

If $x \in \Omega \subset V_0$ then $|x| \leq \mathcal{O}(1)h\ln(1/h)$. Taking $u \in \mathcal{K}er(\tau_N)$ and using Lemma 9 we get

$$\|(h^{-1/2}x)^\alpha(h^{1/2}D_x)^\beta u\|_{\Omega_2} \leq \mathcal{O}\left((\ln(1/h))^{|\alpha|+|\beta|}\right)\|u\|_{\Omega_1\setminus B_{r_0\sqrt{s}}}.$$

Thus the partial differential operator T^r is bounded of norm $\mathcal{O}(h) : H_\Phi(\Omega_1) \mapsto H_\Phi(\Omega_2)$ and as $u \in \mathcal{K}er(\tau_N)$ we have,

$$\|T^r u\|_{\Omega_2} \leq \mathcal{O}(h)\|u\|_{\Omega_1} \leq \mathcal{O}(h)\|u\|_{\Omega_1\setminus B_{r_0\sqrt{s}}}. \tag{82}$$

Then, using (82) with $\Omega_1 = \Omega$ and $\Omega_2 = A^{-1}(\Omega)$, we get

$$\int_{A^{-1}(\Omega)\setminus A^{-1}(B_{r_0\sqrt{s}})} e^{-2\Phi(x)/h}\left|T^r u(x)\right|^2 L(dx) \leq \mathcal{O}(h^2)\int_{\Omega\setminus B_{r_0\sqrt{s}}} e^{-2\Phi(x)/h}\left|u(x)\right|^2 L(dx).$$

We can choose r_0 large enough such that

$$\text{the right hand side of (81)} \leq \frac{1}{4}\int_{\Omega\setminus B_{r_0 e^{-\mu}\sqrt{s}}} e^{-2\Phi(x)/h}|u(x)|^2 L(dx).$$

This implies the following lemma:

Lemma 11. *Suppose $\lambda \in \Lambda_{A,B}$ and $|\lambda| > C$ large enough. Let $h = 1/\mathrm{Re}\,\lambda$, r_0, ϵ be as in Lemma 9. Let Ω be any λ-independent neighborhood of* 0 *and*

$$B_{r_0\sqrt{s}} \subset\subset B_{\epsilon\sqrt{s}} \subset\subset \Omega.$$

Let $A = \mathrm{diag}(e^{\mu_1}, \dots, e^{\mu_n})$ be as in Theorem 3. Let $A^{-1}(\Omega) := \{A^{-1}x,\ x \in \Omega\}$.

If r_0 and ϵ are sufficiently large then for some $m \le 1/2$

$$\int_{\Omega\setminus B_{r_0\sqrt{s}}} e^{-2\Phi/h}|e^{-i2d\lambda}e^{-i\lambda F^r(I;1/\lambda)}u|^2 L(dx) \le m^2 \int_{\Omega\setminus A^{-1}(B_{r_0\sqrt{s}})} e^{-2\Phi/h}|u|^2 L(dx) \tag{83}$$

for any $u \in H_\Phi(\Omega) \cap \mathcal{K}er(\tau_N)$.

10.4. **Bounds useful for the second local Grushin problem, Theorem 7.** We suppose $\lambda \in \Lambda_{A,B}$. Vi fix $N \in \mathbb{N}$ and let $u_N \in H_\Phi(V_0) \cap \mathcal{K}er(\tau_N)$. Let $r_0 > 0$ be large enough. In all lemmas below we use the following convention for the domains:

$$B_{r_0\sqrt{s}} \subset\subset B_{\epsilon_0\sqrt{s}} \subset\subset \Omega_2 \subset\subset \Omega_1 \subset\subset V_0. \tag{84}$$

Lemma 6 implies that the norm of $R_r = M_0 - e^{-i\lambda F^r(I;1/\lambda)}$ is small in $H_\Phi(V_0)$. We have

Lemma 12. *For any $\tilde{N} \in \mathbb{N}$ there is $r = r(N, \tilde{N}, r_0) \in \mathbb{N}$ sufficiently large such that*

$$\|e^{-i2d\lambda}R^r u_N\|^2_{\Omega_2\setminus B_{r\sqrt{s}}} \le \mathcal{O}_r(h^{\tilde{N}})\|u_N\|^2_{\Omega_1\setminus B_{r_0\sqrt{s}}}$$

with domains satisfying (84).

Proof: Let $z = e^{-i2d\lambda}$. We use Lemma 6 (with $\tilde{N}$ instead of N) and Lemma 9:

$$\|zR^r u_N\|^2_{\Omega_2\setminus B_{r_0\sqrt{s}}} \le \mathcal{O}(h^{\tilde{N}})\|u_N\|^2_{\Omega_1} = \mathcal{O}(h^{\tilde{N}})\left(\|u_N\|^2_{B_{r_0\sqrt{s}}} + \|u_N\|^2_{\Omega_1\setminus B_{r_0\sqrt{s}}}\right) \le$$
$$\le \mathcal{O}(h^{\tilde{N}})\left(\|u_N\|^2_{B_{\epsilon\sqrt{s}}\setminus B_{r_0\sqrt{s}}} + \|u_N\|^2_{\Omega_1\setminus B_{r_0\sqrt{s}}}\right) \le \mathcal{O}(h^{\tilde{N}})\|u_N\|^2_{\Omega_1\setminus B_{r_0\sqrt{s}}}.$$

∎

Combining Lemma 11 and Lemma 12 we get

Lemma 13. *Let $M_0 = e^{-i\lambda F^r(I;1/\lambda)} + R^r$. Suppose $\lambda \in \Lambda_{A,B}$, (84) and $A^{-1}(\Omega) = \{A^{-1}x,\ x \in \Omega\}$, $A = \mathrm{diag}(e^{\mu_1}, \dots, e^{\mu_n})$.*

For any $\tilde{N} \in \mathbb{N}$ there is $r \in \mathbb{N}$ such that if r_0 is sufficiently large then for some $m \le 1/2$

$$\|e^{-i2d\lambda}M_0 u_N\|^2_{\Omega_2\setminus B_{r_0\sqrt{s}}} \le m^2\|u_N\|^2_{\Omega_2\setminus A^{-1}(B_{r_0\sqrt{s}})} + \mathcal{O}_r(h^{\tilde{N}})\|u_N\|^2_{\Omega_1\setminus B_{r_0\sqrt{s}}} \tag{85}$$

and

$$\|u_N\|^2_{\Omega_2\setminus B_{r_0\sqrt{s}}} \le \mathcal{O}(1)\|(I - e^{-i2d\lambda}M_0)u_N\|^2_{\Omega_2\setminus B_{r_0\sqrt{s}}} + \mathcal{O}_r(h^{\tilde{N}})\|u_N\|^2_{\Omega_1\setminus B_{r_0\sqrt{s}}}. \tag{86}$$

Proof: We show (86). Choose ϵ such that $\Phi = |x|^2/2$ for $x \in B_{\epsilon\sqrt{s}}$ and $r_0 \le \frac{\sqrt{N+1/2}}{2\sqrt{s\lambda_1}}$ as before. Then, with $z = e^{-i2d\lambda}$, and some constant $m \le 1/2$, bound 85 implies

$$\|(I - zM_0)u_N\|^2_{\Omega_2\setminus B_{r_0\sqrt{s}}} \ge \|u_N\|^2_{\Omega_2\setminus B_{r_0\sqrt{s}}} - \|zM_0u_N\|^2_{\Omega_2\setminus B_{r_0\sqrt{s}}} \ge$$
$$\ge \|u_N\|^2_{\Omega_2\setminus B_{r_0\sqrt{s}}} - m^2\|u_N\|^2_{\Omega_2\setminus A^{-1}(B_{r_0\sqrt{s}})} - \mathcal{O}_r(h^{\tilde{N}})\|u_N\|^2_{\Omega_1\setminus B_{r_0\sqrt{s}}} =$$
$$= (1 - m^2)\|u_N\|^2_{\Omega_2\setminus B_{r_0\sqrt{s}}} - m^2\|u_N\|^2_{B_{r_0\sqrt{s}}\setminus A^{-1}(B_{r_0\sqrt{s}})} - \mathcal{O}_r(h^{\tilde{N}})\|u_N\|^2_{\Omega_1\setminus B_{r_0\sqrt{s}}}. \tag{87}$$

Lemma 9 implies

$$\|u_N\|^2_{B_{r_0\sqrt{s}}\setminus A^{-1}(B_{r_0\sqrt{s}})} \le \|u_N\|^2_{B_{r_0\sqrt{s}}} \le \|u_N\|^2_{B_{\epsilon\sqrt{s}}\setminus B_{r_0\sqrt{s}}} \le \|u_N\|^2_{\Omega_2\setminus B_{r_0\sqrt{s}}}.$$

Then we get that

$$\text{the right hand side of (87)} \geq (1-2m^2)\|u_N\|^2_{\Omega_2\backslash B_{r_0\sqrt{s}}} - \mathcal{O}_r(h^{\tilde{N}})\|u_N\|^2_{\Omega_1\backslash B_{r_0\sqrt{s}}}.$$

Then we have

$$(1-2m^2)\|u_N\|^2_{\Omega_2\backslash B_{r_0\sqrt{s}}} \leq \|(I-zM_0)u_N\|^2_{\Omega_2\backslash B_{r_0\sqrt{s}}} + \mathcal{O}_r(h^{\tilde{N}})\|u_N\|^2_{\Omega_1\backslash B_{r_0\sqrt{s}}}.$$

∎

Lemma 9 implies that $\|u_N\|^2_{B_{r_0\sqrt{s}}} \leq \|u_N\|^2_{\Omega_2\backslash B_{r_0\sqrt{s}}}$ which together with estimate (86) implies

Lemma 14. *Let $\lambda \in \Lambda_{A,B}$. For any $\tilde{N} \in \mathbb{N}$ there is $r \in \mathbb{N}$ such that if r_0 is sufficiently large then*

$$\|u_N\|^2_{\Omega_2} \leq \mathcal{O}(1)\|(I-e^{-i2d\lambda}M_0)u_N\|^2_{\Omega_2} + \mathcal{O}_r(h^{\tilde{N}})\|u_N\|^2_{\Omega_1\backslash B_{r_0\sqrt{s}}} \tag{88}$$

with domains satisfying (84).

11. THE SECOND LOCAL GRUSHIN PROBLEM.

With the same notations as in (74), $M_0 = e^{-i\lambda F^r(I;1/\lambda)} + R^r$, $\lambda \in \Lambda_{A,B}(\epsilon_0)$ verifying (72), for any $\Omega_2 \subset\subset \Omega_1 \subset\subset V_0$, we pose the second Grushin problem:

$$\begin{cases} (I-e^{-i2d\lambda}M_0)u + R_-u_- = v, \\ R_+u = v_+,\ u \in H_\Phi(\Omega_1),\ v \in H_\Phi(\Omega_2),\ u_-, v_+ \in \mathbb{C}^a. \end{cases} \tag{89}$$

Theorem 7. *Let $\lambda \in \Lambda_{A,B}(\epsilon_0)$. We consider the Grushin problem (89).*

Then for any $\tilde{N} \in \mathbb{N}$ there is $r \in \mathbb{N}$ large enough such that we have

(90)

$$\|u\|_{H_\Phi(\Omega_1)} + |u_-|_{\mathbb{C}^a} \leq \mathcal{O}\left((\operatorname{Re}\lambda)^\delta\right)\left(\|v\|_{H_\Phi(\Omega_2)} + |v_+|_{\mathbb{C}^a}\right) + \mathcal{O}_r\left(|\operatorname{Re}\lambda|^{-\tilde{N}}\right)\|u\|_{H_\Phi(\Omega_1)},\ 0<\delta<1.$$

Here δ is as in (69).

Proof: Let $z = e^{-i2d\lambda}$, $h = 1/\operatorname{Re}\lambda$, and N sufficiently large. Applying $(1-\tau_N)$ to the first equation in (89) and using $(1-\tau_N)R_- = 0$, we get $(1-\tau_N)(I-zM_0)u = (1-\tau_N)v$ which implies

$$(I-zM_0)(1-\tau_N)u = (1-\tau_N)v + \tau_N(I-zM_0)(1-\tau_N)u - (1-\tau_N)(I-zM_0)\tau_N u. \tag{91}$$

Then we apply estimate (88) (Lemma 14). Let $B_{r_0\sqrt{s}} \subset\subset \Omega_2 \subset\subset \Omega_1 \subset\subset V_0$. Then

$$\begin{aligned} &\|(1-\tau_N)u\|_{\Omega_2} \leq c\|(1-\tau_N)v\|_{\Omega_2} + \\ &+ \mathcal{O}(1)\|\tau_N(I-zM_0)(1-\tau_N)u - (1-\tau_N)(I-zM_0)\tau_N u\|_{\Omega_2} + \\ &+ \mathcal{O}_r(h^{N_0})\|(1-\tau_N)u\|_{\Omega_1\backslash B_{r_0\sqrt{s}}}. \end{aligned} \tag{92}$$

We need to bound (92).

We have

$$\|\tau_N z e^{-i\lambda F^r(I;1/\lambda)}(1-\tau_N)u\|_{\Omega_2} \leq \mathcal{O}(h^{\tilde{N}})\|u\|_{\tilde{\Omega}_2},$$

where $\Omega_2 \subset\subset \tilde{\Omega}_2 \subset\subset \Omega_1 \subset\subset V_0$, and similar, interchanging τ_N and $1-\tau_N$. It follows from the following facts:

$$\text{if } u \in O_N = \mathcal{K}er\,\tau_N,\ \text{then } u^{(\beta)}(e^{-\mu}x)(e^{-\mu}x)^\beta \in O_N\ \forall\beta,$$

and formula (80), which implies that $\tau_N z e^{-i\lambda F^r(I;1/\lambda)}(1-\tau_N) = 0$ for any fixed order r.

Then it is enough to estimate $\|\tau_N zR(1-\tau_N)\|_{\Omega_2}$ and $\|(1-\tau_N)zR\tau_N\|_{\Omega_2}$ which can be done directly, but from more general result, Lemma 6 on V_0, we know that for any $\tilde{N}$ we can find r such that $(92) \leq \mathcal{O}_r(h^{\tilde{N}})\|u\|_{\tilde{\Omega}_2}$.

We get the estimate

$$\|(1-\tau_N)u\|_{\Omega_2} \le \mathcal{O}(1)\|(1-\tau_N)v\|_{\Omega_2} + \mathcal{O}_r(h^{\tilde N})\|u\|_{\tilde\Omega_2}. \tag{93}$$

Apply τ_N to (89). Then

$$\begin{cases} \tau_N(I - zM_0)u + R_-u_- & = \tau_N v \\ R_+\tau_N u & = v_+ \end{cases}$$

since $\tau_N R_- = R_-$ and $R_+ = R_+\tau_N$. Then

$$\begin{cases} \tau_N(I - ze^{-i\lambda F^r(I;1/\lambda)})\tau_N u + R_-u_- & = \tau_N v - \tau_N(I - zM_0)(1-\tau_N)u - \\ & \quad \tau_N z\underbrace{(e^{-i\lambda F^r(I;1/\lambda)} - M_0)}_{R^r}\tau_N u \\ R_+\tau_N u & = v_+. \end{cases}$$

The first term in $-\tau_N(I - zM_0)(1-\tau_N)u - \tau_N zR^r\tau_N u$ which is equal to $\tau_N zM_0(1-\tau_N)u = \tau_N ze^{-i\lambda F^r(I;1/\lambda)}(1-\tau_N)u + \tau_N zR^r(1-\tau_N)u$ is already estimated. The second term can be bounded as before using Lemma 6 in Section 6.

Then estimate (75) implies

$$\|\tau_N u\|_{\Omega_1} + |u_-|_{\mathbb{C}^a} \le \mathcal{O}\left(h^{-\delta}\right)\left(\|\tau_N v\|_{\Omega_2} + |v_+|_{\mathbb{C}^a}\right) + \mathcal{O}_r(h^{\tilde N})\|u\|_{\tilde\Omega_2}. \tag{94}$$

Adding (93) and (94) we get

$$\|u\|_{\Omega_1} + |u_-|_{\mathbb{C}^a} \le \mathcal{O}(1)\|(1-\tau_N)v\|_{\Omega_2} + \mathcal{O}\left(h^{-\delta}\right)\left(\|\tau_N v\|_{\Omega_2} + |v_+|_{\mathbb{C}^a}\right) + \mathcal{O}_r(h^{\tilde N})\|u\|_{\tilde\Omega_2}. \tag{95}$$

Then using Proposition 4.3 in [8], we get the desired estimate (90). ∎

12. THE GLOBAL GRUSHIN PROBLEM

We denote $\Omega_{\text{int}} \in T^*\partial\Omega_1$ the λ-independent neighborhood of a_1 such that $W_0 \subset\subset \Omega_{\text{int}} \subset\subset W_1$, where W_0, W_1 are defined in (22), (26). Let Ω_{ext} be such that $\complement\Omega_{\text{ext}} \subset\subset \Omega_{\text{int}}$.

We consider the original operator $I - M(\lambda)$ in $\mathcal{H}$ equipped with the norm

$$|u|_{\mathcal{H}} := \|(1-\chi_2)T_{\Lambda_{tG}}u\|_{L^2_{tG}(T^*(\partial\Omega_1))} + \|\chi_1 Uu\|_{L^2(e^{-2\Phi/h}L(dx))} =: |u|_{\Omega_{\text{ext}}} + |u|_{\Omega_{\text{int}}},\ u \in \mathcal{H},$$

where $\chi_1 \in C_0^\infty(\mathbb{C}^n)$ is equal to 1 in a neighborhood of $a_1 = 0$, with the support independent of λ and $W_0 \subset\subset \operatorname{supp}\chi_1 \subset\subset \Omega_{\text{int}} \subset\subset W_1$, and $\chi_2 \in C_0(T^*X)$ equal to 1 near $(a_1, 0)$ and essentially the same function as χ_1 after suitable identification of domains. Here $L^2_{tG}(T^*(\partial\Omega_1))$ stands for $L^2(\Lambda; e^{-2H/h}|\langle\alpha_\xi\rangle|^{2m}d\alpha)$. Let

$$U:\ \mathcal{H} \mapsto H_\Phi(\Omega_{\text{int}}),\ U = B_rFT_{\text{Bargman}},\ V_1 \supset \text{neigh}(a_1) \mapsto \text{neigh}(0) \subset \hat V_1,$$

$\pi(W_1) = V_1$, be as before (we omit $\hat{}$) such that

$$UM(\lambda)u = e^{-2id\lambda}M_0Uu + \mathcal{O}\left(h^N\right)\|u\|_{\mathcal{H}},\ \text{in } H_\Phi(\Omega_{\text{int}}),\ M_0 - e^{-i\lambda F^r} \equiv \mathcal{O}(\rho^{2r+1}, h^{2r+1}),$$

where $h = 1/\operatorname{Re}\lambda,\ \lambda \in \Lambda_{A,B}$.

Let $V:\ H_\Phi(\Omega_{\text{int}}) \mapsto H(\Lambda_{tG})$ be the approximate microlocal inverse of U such that if $\chi(x, hD_x)$ is pseudodifferential operator adapted to $H(\Lambda_{tG})$ with compact symbol and with $\operatorname{supp}\chi \subset\subset \Omega_{\text{int}}$, then $\chi(VU - I)$ and $(UV - I)\chi$ are neglectible, $VUu - u = \mathcal{O}(h^N)\|u\|_{\mathcal{H}}$.

Theorem 8. *For a given $(v, v_+) \in \mathcal{H}\times\mathbb{C}^a$ consider the Grushin problem*

$$\begin{cases} (I - M)u + VR_-u_- & = v, \\ R_+Uu & = v_+. \end{cases} \tag{96}$$

For $\lambda \in \Lambda_{A,B}(\epsilon_0)$*, verifying (72), and* $\operatorname{Re}\lambda$ *sufficiently large the Grushin problem (96) has a unique solution* $(u, u_-) \in \mathcal{H} \times \mathbb{C}^a$*. Moreover, we have the a priori estimate*

$$|u|_{\mathcal{H}} + |u_-|_{\mathbb{C}^a} \le \mathcal{O}((\operatorname{Re}\lambda)^{\delta})\left(|v|_{\mathcal{H}} + |v_+|_{\mathbb{C}^a}\right) \tag{97}$$

with some δ*,* $0 < \delta < 1$*.*

Proof: The existence of a solution follows as in [9]. We need to show estimate (97). We apply U from the left in the first equation in (96) and denote $Uu = \tilde{u}$. Then

$$\begin{cases} (I - e^{-2id\lambda}M_0)\tilde{u} + R_-u_- &= Uv + \tilde{w},\ \tilde{w} := (UM - e^{-2id\lambda}M_0U)u + (I - UV)R_-u_-, \\ R_+\tilde{u} &= v_+. \end{cases}$$

Denote $\tilde{v} = Uv + \tilde{w}$. Introduce as before Ω_{ext}, Ω_{int}, W_0, $V_0 = \pi(W_0)$. Let

$$\Omega_2 \subset\subset \Omega_1 \subset\subset V_0 \subset\subset \Omega_3' \subset\subset \Omega_2' \subset\subset \Omega_1' \subset\subset \Omega_{\text{int}}.$$

Estimate on V_0 **:**
By (90) we have:

$$\|\tilde{u}\|_{H_\Phi(\Omega_1)} + |u_-|_{\mathbb{C}^a} \le \mathcal{O}\left(h^{-\delta}\right)\left(\|\tilde{v}\|_{H_\Phi(\Omega_2)} + |v_+|_{\mathbb{C}^a}\right) + \mathcal{O}_r\left(h^N\right)\|\tilde{u}\|_{H_\Phi(\Omega_1)},\ 0 < \delta < 1. \tag{98}$$

We have

$$\|\tilde{w}\|_{H_\Phi(\Omega_2)} \le \mathcal{O}_r(h^N)\left(\|\tilde{u}\|_{H_\Phi(\Omega_1)} + |u_-|_{\mathbb{C}^a}\right). \tag{99}$$

This gives

$$\|\tilde{u}\|_{H_\Phi(\Omega_1)} + |u_-|_{\mathbb{C}^a} \le \mathcal{O}\left(h^{-\delta}\right)\left(\|Uv\|_{H_\Phi(\Omega_2)} + |v_+|_{\mathbb{C}^a}\right) + \mathcal{O}_r\left(h^N\right)\left(\|\tilde{u}\|_{H_\Phi(\Omega_1)} + |u_-|_{\mathbb{C}^a}\right). \tag{100}$$

Estimate on $\Omega_{\text{int}} \setminus V_0$ **:**
In $(I - zM_0)\tilde{u} = -R_-u_- + \tilde{v}$ we use (a variant of) Theorem 3, estimate (50), which implies that there is $N_1 > 0$ such that

$$\|z\chi M_0\tilde{u}\|_{L^2_\Phi(\Omega_2')} \le \mathcal{O}(h^{N_1})\|\tilde{u}\|_{H_\Phi(\Omega_1')},\ \operatorname{supp}\chi \subset \Omega_{\text{int}} \setminus V_0.$$

We have then

$$\|\chi(1 - zM_0)\tilde{u}\|_{L^2_\Phi(\Omega_2')} \ge \|\chi\tilde{u}\|_{L^2_\Phi(\Omega_1')} - \mathcal{O}(h^{N_1})\|\tilde{u}\|_{H_\Phi(\Omega_1')}$$

and

$$\|\chi\tilde{u}\|_{L^2_\Phi(\Omega_1')} \le \|\chi R_-u_-\|_{L^2_\Phi(\Omega_2')} + \|\chi\tilde{v}\|_{L^2_\Phi(\Omega_2')} + \mathcal{O}(h^{N_1})\|\tilde{u}\|_{H_\Phi(\Omega_1')}.$$

This implies for any $N \in \mathbb{N}$

$$\|\chi\tilde{u}\|_{L^2_\Phi(\Omega_1')} \le \mathcal{O}(h^N)|u_-|_{\mathbb{C}^a} + \|\chi\tilde{v}\|_{L^2_\Phi(\Omega_2')} + \mathcal{O}(h^{N_1})\|\tilde{u}\|_{H_\Phi(\Omega_1')}. \tag{101}$$

Similar, applying (50) to each term in $\tilde{w}$ separately, we get

$$\|\chi\tilde{w}\|_{L^2_\Phi(\Omega_2')} \le \mathcal{O}(h^{N_1})\left(\|\chi\tilde{u}\|_{L^2_\Phi(\Omega_1')} + |u_-|_{\mathbb{C}^a}\right) \tag{102}$$

and we have

$$\|\chi\tilde{u}\|_{L^2_\Phi(\Omega_1')} \le \mathcal{O}(h^{N_1})|u_-|_{\mathbb{C}^a} + \|\chi Uv\|_{L^2_\Phi(\Omega_2')} + \mathcal{O}(h^{N_1})\|\tilde{u}\|_{H_\Phi(\Omega_1')}. \tag{103}$$

Applying (100) and (103) with $\tilde{u}$ replaced by Uu we get

$$|u|_{\Omega_1'} + |u_-|_{\mathbb{C}^a} \le \mathcal{O}(h^{-\delta})\left(|v|_{\Omega_2'} + |v_+|_{\mathbb{C}^a}\right) + \mathcal{O}\left(h^{N_1}\right)\left(|u|_{\Omega_1'} + |u_-|_{\mathbb{C}^a}\right), \tag{104}$$

where $\Omega_2' \subset\subset \Omega_1' \subset\subset \Omega_{\text{int}}$.
Estimate on Ω_{ext} **:**
In order to estimate $u = Mu - VR_-u_- + v$ we use bound (42) with some $N_2 > 0$:

$$|Mu|_{\Omega_{\text{ext}}} \le \mathcal{O}(h^{N_2})|u|_{\Omega_{\text{ext}}}$$

and get for any $N \in \mathbb{N}$

$$|u|_{\Omega_{\text{ext}}} \leq |Mu|_{\Omega_{\text{ext}}} + |v|_{\Omega_{\text{ext}}} + |VR_-u_-|_{\Omega_{\text{ext}}} \leq \mathcal{O}(h^{N_2})|u|_{\Omega_{\text{ext}}} + |v|_{\Omega_{\text{ext}}} + \mathcal{O}(h^N)|u_-|.$$

Summing up we get

$$\begin{aligned} &|u|_{\Omega_{\text{ext}}} + |u|_{\Omega_{\text{int}}} + |u_-|_{\mathbb{C}^a} \leq \\ &\leq \mathcal{O}(h^{-\delta})\left(|v|_{\Omega_{\text{ext}}} + |v|_{\Omega_{\text{int}}} + |v_+|_{\mathbb{C}^a}\right) + \mathcal{O}(h^{N_0})\left(|u|_{\Omega_{\text{ext}}} + |u|_{\Omega_{\text{int}}} + |u_-|_{\mathbb{C}^a}\right), \end{aligned}$$

with $N_0 = \min(N, N_1, N_2)$. The last term can be absorbed in the left hand side and we get estimate (97).

Then it is clear that

$$\mathcal{P} = \begin{pmatrix} I - M(\lambda) & VR_- \\ R_+U & \end{pmatrix} : \ \mathcal{H} \times \mathbb{C}^a \mapsto \mathcal{H} \times \mathbb{C}^a$$

is Fredholm of index 0. Estimate (97) implies that $\mathcal{P}$ is injective and thus bijective for $\lambda \in \Lambda_{A,B}(\epsilon_0)$.

∎

Denote

$$\mathcal{E} = \begin{pmatrix} E & E_+ \\ E_- & E_{+-} \end{pmatrix}$$

the inverse of $\mathcal{P}$. Let $h = 1/\operatorname{Re}\lambda$, $\lambda \in \Lambda_{A,B}$. Then it is known

$$0 \in \sigma(I - M(\lambda)) \ \Leftrightarrow \ 0 \in \sigma(E_{+-}(\lambda)).$$

Let $v_+ \in \mathbb{C}^a$. If $(u, u_-) = \mathcal{E}(0, v_+)$ then $u = E_+(v_+)$, $u_- = E_{+-}(v_+)$.

Let

$$\begin{cases} (I - M)u + VR_-u_- &= 0, \\ R_+Uu &= v_+. \end{cases}$$

Then we have

$$(I - e^{-2id\lambda}M_0)\tilde{u} + R_-u_- = \tilde{w} = (UM - e^{-2id\lambda}M_0U)u + (I - UV)R_-u_-$$

and $\tilde{w}$ satisfies (99):

$$\|\tilde{w}\|_{H_\Phi(\Omega)} \leq \mathcal{O}_r(h^N)\left(|u|_{\mathcal{H}} + |u_-|_{\mathbb{C}^a}\right) = \mathcal{O}_r(h^N)|v_+|_{\mathbb{C}^a}.$$

We have then for r large enough

$$\begin{cases} \tau_J\left(I - e^{-2id\lambda}e^{-i\lambda F^r(I(1/\lambda);1/\lambda)}\right)\tau_J\tilde{u} + R_-u_- &= \mathcal{O}_r(h^N)|v_+|_{\mathbb{C}^a}, \\ \tau_J\tilde{u} &= R_-v_+ \end{cases}$$

and

$$\begin{aligned} \left(I - e^{-2id\lambda}e^{-i\lambda F^r(I(1/\lambda);1/\lambda)}\right)R_-v_+ + R_-u_- = \mathcal{O}_r(h^N)|v_+|_{\mathbb{C}^a} \ \Leftrightarrow \\ u_- = -R_+\left(I - e^{-2id\lambda}e^{-i\lambda F^r(I(1/\lambda);1/\lambda)}\right)R_-v_+ + \mathcal{O}_r(h^N)|v_+|_{\mathbb{C}^a}. \end{aligned}$$

We have then for $\lambda \in \Lambda_{A,B}(\epsilon_0)$:

$$E_{+-}(\lambda) = E^0_{+-}(\lambda) + \mathcal{O}_r\left((\operatorname{Re}\lambda^r_{\alpha,k})^{-N}\right), \text{ where } E^0_{+-} = -\left\{\left(1 - e^{i2d\lambda}K_\alpha(\lambda)\right)\right\}_{|\alpha\in J}.$$

Let $\gamma = \gamma(\lambda^r_{\alpha,k})$ be simple loop such that $E^0_{-+}(\lambda)$ is invertible for $\lambda \in \gamma$. Suppose

$$\lambda \in \gamma \ \leftrightarrow \ \lambda = \lambda^r_\alpha + \mathcal{O}_r\left((\operatorname{Re}\lambda^r_{\alpha,k})^{-N}\right),$$

where $\lambda^r_{\alpha,k}$ is solution of (68). Then the number of poles of $(I-M(\lambda))^{-1}$ inside γ counted with multiplicity is equal to the number of roots $m(\gamma)=m(\lambda^r_{\alpha,k})$ inside γ of $\det E^0_{-+}(\lambda)=0$,

$$m(\gamma)=\frac{1}{2\pi i}\int_\gamma \frac{\left(\det E^0_{-+}(\lambda)\right)'}{\det E^0_{-+}}d\lambda.$$

We have $0\in\sigma(I-M(\lambda))$ if $\lambda=\lambda^r_\alpha+\mathcal{O}_r\left((\operatorname{Re}\lambda^r_{\alpha,k})^{-N}\right)$.

This accomplishes the proof of Theorem 2. ▮

Appendix A. Some facts about the quantum billiard operator M.

A.1. **Boundness.** Let Ω_1 be an obstacle with analytic boundary, non-trapping. Let $\Omega_1\subset B(0,R)$. Denote

$$U_{A,B}:=\{\lambda;\ \operatorname{Im}\lambda\le A|\lambda|^{1/3}-B\}. \tag{105}$$

Let $R(\lambda)$ be the outgoing Dirichlet resolvent in $\complement\Omega_1$.

Theorem 9. *Let $\chi\in C_0^\infty(\mathbb{R}^n)$ be such that $\chi\equiv 1$ in $B(0,R)$. There exist $A,B>0$ such that $\chi R\chi(\lambda)$, defined for $\operatorname{Im}\lambda<0$ has an analytic extension to the domain (105) as a bounded operator $L^2(\complement\Omega_1)\mapsto H^1_0(\complement\Omega_1)$, satisfying the estimate*

$$\exists C,D>0;\ \|\chi R\chi\|_{\mathcal{L}(L^2(\complement\Omega_1),H^1_0(\complement\Omega_1))}\le Ce^{D\operatorname{Im}\lambda^+},$$

where $\operatorname{Im}\lambda^+=\max(\operatorname{Im}\lambda,0)$.

For $i=1,2$ let

$$H_{i,+}(\lambda):\ C^\infty(\partial\Omega_i)\mapsto C^\infty(\overline{\mathbb{R}^{n+1}\setminus\Omega_i})$$

be the outgoing resolvent of the problem

$$\begin{cases}(\Delta+\lambda^2)H_{i,+}(\lambda)u=0 \text{ in } \mathbb{R}^{n+1}\setminus\Omega_i\\ H_{i,+}(\lambda)u_{|\partial\Omega_i}=u\end{cases}$$

extended as an operator $H^{1/2}(\partial\Omega_i)\mapsto H^1_{\mathrm{loc}}(\overline{\complement\Omega_i})$. In [2] it is proven that this resolvent, analytical for $\operatorname{Im}\lambda<0$ has an analytical extension to the domain of the form (105) as a bounded operator $H^{1/2}(\partial\Omega_i)\mapsto H^1_{0,\mathrm{loc}}(\overline{\complement\Omega_i})$ and satisfy the following estimate:

$$\forall R>0,\ \exists C>0,\ \exists D>0,\ \forall\lambda\in U_{A,B},\ \|H_{i,+}\|_{\mathcal{L}(H^{1/2}(\partial\Omega_1),H^1(\complement\Omega_1\cap B(0,R)))}\le Ce^{D\operatorname{Im}\lambda^+}.$$

We denote $H_{21}(\lambda)=H_{2,+}\circ\gamma_2\circ H_{1,+}$, where γ_i is the operator of restriction to $\partial\Omega_i$, the outgoing resolvent of the problem

$$\begin{cases}(\Delta+\lambda^2)H_{21}u &=0, \text{ in } \complement\Omega_2\\ \qquad H_{21}u_{|\partial\Omega_2} &=H_{1,+}u_{|\partial\Omega_2}.\end{cases}$$

We define $H_i(\lambda)u=H_{i,+}(\lambda)u_{|\partial\Omega_{i+1}}$, where $\partial\Omega_3=\partial\Omega_1$ and

$$M(\lambda)=H_2(\lambda)H_1(\lambda)=\gamma_1H_{2,+}\gamma_2H_{1,+}.$$

Lemma 15 (Burq)**.** *Operator $M(\lambda)$ defined on $H^1(\partial\Omega_1)\mapsto H^1(\partial\Omega_1)$ for $\operatorname{Im}\lambda<0$ has an analytic extension in the domain of the form (105) and there satisfies the following estimate*

$$\exists D>0,\ \exists C>0,\ \forall\lambda\in U_{A,B},\ \|M(\lambda)\|_{\mathcal{L}(H^{1/2}(\partial\Omega_1))}\le C|\lambda|^2e^{D\operatorname{Im}\lambda^+}$$

Moreover, as in a neighborhood of Ω_1, Mu satisfies $(\Delta+\lambda^2)Mu=0$, we get

$$\exists D>0,\ \forall s\in\mathbb{R},\ \exists C_s>0,\ \forall\lambda\in U_{A,B},\ \|M(\lambda)\|_{\mathcal{L}(H^s(\partial\Omega_1))}\le C_se^{D\operatorname{Im}\lambda^+}|\lambda|^{s+2}.$$

A.2. **Microlocal unitarity of M with respect to the flux norm.** We suppose that λ is real. Denote $h = 1/\lambda$. Let v satisfy

$$\left\{ \begin{aligned} Pu := (-h^2\Delta - 1)u &= 0, \ u \text{ outgoing} \\ \gamma_1 u &= v. \end{aligned} \right. \tag{106}$$

We define $H_{1,+} : \ \mathcal{D}'(\partial\Omega_1) \mapsto \mathcal{D}'(\overline{\complement\Omega_1})$ the solution operator. In the similar way we define $H_{2,+} : \ \mathcal{D}'(\partial\Omega_2) \mapsto \mathcal{D}'(\overline{\complement\Omega_2})$, which satisfies $PH_{2,+}v = 0$ and $\gamma_2 H_{2,+}v = v$. Let $H_1 = \gamma_2 H_{1,+}$ and $H_2 = \gamma_1 H_{2,+}$. Then $M = H_2 H_1 : \ H^1(\partial\Omega_1) \mapsto H^1(\partial\Omega_1)$.

The billiard operator M can be identified with the monodromy operator M as in [24] in the form presented in ([14], p. 360). Let $\ker_{\rho_1}(P)$ be local kernel P near $\rho_1 = (a_1, 0)$. The solution operator K in ([14]) is the operator $H_{1,+}$,

$$H_{1,+}v \in \ker_{\rho_1}(P), \ H_{1,+}v_{|\partial\Omega_1} = v.$$

We have

$$\mathcal{M} = H_{1,+}\gamma_1 H_{2,+}\gamma_2 : \ \ker_{\rho_1}(P) \mapsto \ker_{\rho_1}(P).$$

As in ([14]) we have identification $\ker_{\rho_1}(P) \simeq \mathcal{D}'(\partial\Omega_1)$ via $K = H_{1,+}$. Then the monodromy operator M on $\mathcal{D}'(\partial\Omega_1)$ satisfies

$$KMv = \mathcal{M}Kv, \ v \in \mathcal{D}'(\partial\Omega_1).$$

Let $\chi \in C^\infty(\overline{\complement\Omega_1})$ be a microlocal cut-off function supported in a neighborhood W_2 of Ω_1 such that $\chi = 1$ in $\complement\Omega_1 \cap W_1$, where $W_1 \subset\subset W_2$.

We define the *quantum flux* norm (see [14], p. 360) on the outgoing solutions u of (106) as follows

$$\|u\|^2_{\mathrm{QF}} := \langle \frac{i}{h}[P, 1-\chi]u|u\rangle.$$

It is easy to see that $\|u\|_{\mathrm{QF}}$ is independent of χ which implies that M is microlocally unitary with respect to $\|.\|_{\mathrm{QF}}$ for real h :

Lemma 16. *The billiard operator M is microlocally unitary for all real λ with respect to $\|.\|_{\mathrm{QF}}$: $\|\mathcal{M}Kv\|_{\mathrm{QF}} = \|H_{1,+}Mv\|_{\mathrm{QF}} = \|H_{1,+}v\|_{\mathrm{QF}} + \mathcal{O}(h^\infty)$, where $u = H_{1,+}v$ is the outgoing solution of (106) and $WF_h(v) \subset \mathrm{neigh}(\rho_1)$. Here the wave front set $WF_h(v)$ is defined as in* [9].

Using the Green's formula in a neighborhood of Ω_1 bounded by $\partial\Omega_1$ on one side, we get

$$\begin{aligned} \|u\|^2_{\mathrm{QF}} &= \langle \frac{i}{h}[P, 1-\chi]u|u\rangle = \frac{i}{h}\langle (h^2\Delta + 1)\chi u - \chi(h^2\Delta+1)u|u\rangle = \\ &= \frac{h}{i}\int_{\partial\Omega_1} \{(\partial_\nu H_{1,+}v)\overline{H_{1,+}v} - H_{1,+}v(\partial_\nu \overline{H_{1,+}v})\} S(dx) = \\ &= 2\mathrm{Re}\,\langle \frac{h}{i}\partial_\nu H_{1,+}v|v\rangle_{L^2(\partial\Omega_1)} = 2\mathrm{Re}\,\langle Av|v\rangle_{L^2(\partial\Omega_1)}, \ u = H_{1,+}v, \end{aligned}$$

where ∂_ν is the normal derivative, $Av := \gamma_1 \frac{h}{i}\partial_\nu H_{1,+}v$. As A is pseudodifferential operator of order 0, elliptic near $(a_1, 0)$ (we use the explicit WKB construction of $H_{1,+}$ in the hyperbolic zone) then we have for v with $WF_h(v) \subset \mathrm{neigh}(\rho_1)$

$$\|H_{1,+}v\|_{\mathrm{QF}} \sim \|v\|_{L^2(\partial\Omega_1 \cap \mathrm{neigh}(\rho_1))}.$$

Lemma 17. *There exists pseudodifferential operator B of order 0 elliptic near a_1 such that BMB^{-1} is microlocally unitary for all real λ with respect to $\|.\|_{L^2(\partial\Omega_1)}$:*

$$\|BMB^{-1}v\|_{L^2(\partial\Omega_1)} = \|v\|_{L^2(\partial\Omega_1)} + \mathcal{O}(h^\infty), \ WF(v) \subset \mathrm{neigh}(a_1, 0).$$

References

[1] V.M. Babich and N.S. Grigoreva. The analytic continuation of the resolvent of the exterior three dimensional problem for the Laplace operator to second sheet. *Funktsional. Anal. i Prilozhen.*, 1(8):71–74, 1974.

[2] C. Bardos and G. Lebeau and J. Rauch. Scattering frequencies and Gevrey 3 singularities. *Inv. Math.*, 90(1):77–114, 1987.

[3] C. Bardos and J.C. Guillot and J. Ralston. La relation de Poisson pour l'équation des ondes dans un ouvert non borné. application à la théory de la diffusion. *C.P.D.E.*, 7(8):905–958, 1982.

[4] N. Burq. Pôles de diffusion engendrés par un coin. *Astérisque*, 242(6), 1997.

[5] N. Kaidi and P. Kerdelhué. Forme normale de Birkhoff et résonances. *Asymptotic Analysis*, (23):1–21, 2000.

[6] J.-P. Françoise. Propriétés de généricité des transformations canoniques. In J. Palis Jr, editor, *Geometric dynamics Proceedings, Rio de Janeiro, 1981, Lecture Notes in Math.*, number 1007, pages 216–260. Springer, 1983.

[7] T. Hargé and G. Lebeau. Diffraction par un convexe. *Invent. Math.*, 1(118):161–196, 1994.

[8] C. Gérard and J. Sjöstrand. Semi-classical resonances generated by a closed trajectory of hyperbolic type. *Commun. Math. Phys.*, 108:391–421, 1987.

[9] Christian Gérard. Asymptotique de pôles de la matrice de scattering pour deux obstacles strictement convexes. *Supplément au Bulletin de la Société Mathématique de France*, 1988.

[10] B. Helffer and J. Sjöstrand. Résonances en limite semi-classique. *Bulletin de la Société de France, Mémoire (nouvelle série)*, 24/25:1–228, 1986.

[11] L. Hörmander. *The analysis of linear partial differenrial operators*, volume I–IV. Springer Verlag, 1985.

[12] A. Iantchenko. La forme normale de Birkhoff pour un opérateur intégral de Fourier. *Asymptotic Analysis*, 17(1):71–92, 1998.

[13] A. Iantchenko and J. Sjöstrand. Birkhoff normal forms for Fourier integral operators II. *American Journal of Mathematics*, 124:817–850, 2002.

[14] A. Iantchenko and J. Sjöstrand and M. Zworski. Birkhoff normal forms in semi-classical inverse problems. *Mathematical Research Letters*, 9:337–362, 2002.

[15] M.Ikawa. On the poles of scattering matrix for two convex obstacles. *J. Math. Kyoto Univ.*, 23:127–194, 1983.

[16] Gille Lebeau. Régularité Gevrey 3 pour la diffraction. *Comm. P.D.E*, 9:1437–1494, 1984.

[17] V. Petkov. Exposé au Séminaire Goulaouic-Schwartz 82–83.

[18] V. Petkov and L. Stoyanov. *Geometry of Reflected Rays and Inverse Spectral Problems.* John Wiley & Sons, New York, 1992.

[19] G. Popov. Some estimates of Green's functions in the shadow. *Osaka J. Math.*, 24(1):1–12, 1987.

[20] J. Sjöstrand and M. Zworski. Complex scaling and the distribution of scattering poles. *J. Amer. Math. Soc.*, (4):729–769, 1991.

[21] J. Sjöstrand and M. Zworski. Estimates on the number of scattering poles near the real axis for strictly convex obstacles. *Ann. Inst. Fourier*, 3(43):769–790, 1993.

[22] J. Sjöstrand. *Singularités Analytiques Microlocales*, volume 95. Astérisque, 1982.

[23] J. Sjöstrand. Density of resonances for strictly convex analytic obstacles. *Can. J. Math.*, 48(2):397–447, 1996.

[24] J. Sjöstrand and M. Zworski. Quantum monodromy and semi-classical trace formulæ. to appear in J. Math. Pure Appl., published in `http://xxx.lanl.gov/abs/math.SP/0108052`.

[25] Johannes Sjöstrand. Semi-classical resonances generated by non degenerate critical points. In *Collection Pseudo-Diff. Operators, Lecture Notes in Math.*, volume 1256, pages 402–429. Oberwolfach, 1986.

[26] Johannes Sjöstrand and Maciej Zworski. Asymptotic distribution of resonances for convex obstacles. *Acta Math.*, 183(2):191–253, 2000.

[27] Shlomo Sternberg. Infinite Lie groups and the formal aspects of Dynamics. *Journal of Math. and Mechanics*, 10(3):451–476, 1961.

[28] Counting scattering poles M. Zworski. *Spectral and Scattering Theory*, pages 301–331. (M. Ikawa, ed), Marcel Dekker, New York, 1994.

[29] M. Zworski. Dimension of the limit set and the density of resonances for convex co-compact hyperbolic surfaces. *Inventiones Math.*, 136:353–409, 1999.

Malmö University, School of Technology and Society, SE-205 06 Malmö, Sweden
E-mail address: `ai@ts.mah.se`